

Polyclonal Antibody to Biotin - Texas Red

Catalog No.:	R1615TR
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Background:	Biotin is a water soluble vitamin, generally classified as a B complex vitamin, also called vitamin B4. After the initial discovery of biotin, nearly forty years of research were required to establish it as a vitamin. Biotin is required by all organisms but can only be synthesized by bacteria, yeasts, molds, algae, and some plant species. Biotin is required as prosthetic group of enzymes involved in incorporation of carbon dioxide into organic compounds. Biotin has a MW of 244 Da.
Host:	Goat
Immunogen:	Biotin conjugated to Keyhole Limpet Hemocyanin (KLH)
Format:	State: Lyophilized purified IgG fraction Purification: Immunoaffinity chromatography Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative Label: Texas Red – - Sulfonyl Chloride (TR; Molecular Weight 625 daltons) <i>Absorption / Emission:</i> 596 nm / 620 nm <i>Molar Ratio:</i> 4.8 moles Texas Red™ per mole of Goat IgG F(ab') ₂ . Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Flow cytometry: 1:2,000 - 1:10,000. IF microscopy: 1:500 - 1:2,500. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Biotin coupled to sepharose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, biotinylated bovine serum albumin and biotinylated IgG.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage:

Store vial at 2-8°C prior to restoration.

Restore with 1.0 ml of deionized water (or equivalent). Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Dilute only prior to immediate use.

Avoid cycles of freezing and thawing.

Shelf life: one year from despatch.

General References: 1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com