

Polyclonal Antibody to Chicken IgG [H&L] - Biotin-

Alternate names:	Chicken IgY
Catalog No.:	R1617B
Quantity:	2 mg
Concentration:	2.0 mg/ml (before lyophil. by UV absorbance at 280 nm)
Host:	Goat
Immunogen:	Chicken IgG whole molecule.
Format:	State: Lyophilized purified IgG fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer. Label: Biotin – Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) <i>Molar Ratio:</i> 10-20 BAC molecules per Goat IgG molecule Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. This product has been assayed against 1.0 ug of Chicken IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:80,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Chicken IgG and Chicken Serum.
Storage:	Store vial at 2-8°C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: one year from despatch.

General References: 1. Bayer & Wilchek Methods in Enzymology 184; 138-160, 1990.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com