

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com R1617TR Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com



Polyclonal Antibody to Chicken IgG [H&L] -Texas Red-

Alternate names:	Chicken IgY
Catalog No.:	R1617TR
Quantity:	2 mg
Concentration:	2.0 mg/ml (by UV absorbance at 280 nm)
Host:	Goat
Immunogen:	Chicken IgG whole molecule.
Format:	 State: Lyophilized purified IgG fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Texas Red – - Sulfonyl Chloride (Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 4.7 moles Texas RedTM per mole of Goat IgG. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Imunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Chicken IgG and Chicken Serum.
Storage:	Store vial at 2-8°C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: one year from despatch.

General References: 1. J.A. Titus, P.P. Haugland, S.D. Sharrow, D.M. Segal J. Immunol. Methods 50; 193, 1982.