

Monoclonal Antibody to CD204 - FITC

Alternate names:	MSR1, Macrophage acetylated LDL receptor I and II, Macrophage scavenger receptor types I and II, SCARA1, Scavenger receptor class A member 1
Catalog No.:	SM029FT
Quantity:	25 µg
Concentration:	0.1 mg/ml
Background:	CD204 is expressed by tissue macrophages and functions both as an endocytic receptor for lipoproteins and as an adhesion receptor for macrophages binding to ligand rich tissues e.g. atherosclerotic lesions. Clone 2F8 inhibits the uptake of acetylated low-density lipoproteins and also inhibits divalent cation independent adhesion. Recent research shows that clone 2F8 recognises an epitope within SRA that is polymorphic in the SRA from C57BL/6 mice. Clone 2F8 is therefore unsuitable for use with the C57BL/6 mouse strain.
Uniprot ID:	P30204
NCBI:	NP_001106797.1
GeneID:	20288
Host / Isotype:	Rat / IgG2b
Clone:	2F8
Immunogen:	RAW264 cell line. Remarks: Spleen cells from immunised AO rats were fused with cells of the Y3 rat myeloma cell line.
Format:	State: Liquid purified IgG fraction. Purification: Affinity Chromatography on Protein G. Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer. Label: FITC – Fluorescein Isothiocyanate Isomer 1
Applications:	Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells in 100 µl. Use The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises the Murine Scavenger Receptor class A (SRA), type I and II, also known as CD204. Species: Mouse. Reacts with Pig. Other species not tested.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light. Shelf life: one year from despatch.

General References:

1. Fraser, I.P. et al. (1993) Divalent cation-independent macrophage adhesion inhibited by monoclonal antibody to murine scavenger receptor. *Nature* 364: 343-346.
2. de Villiers, W.J.S. et al. (1994) Macrophage-colony-stimulating factor selectively enhances macrophage scavenger receptor expression and function *J. Exp. Med.* 180: 705-709.
3. Hughes, D.A. et al. (1995) Murine Macrophage Scavenger Receptor: in vivo expression and function as receptor for macrophage adhesion in lymphoid and non-lymphoid organs. *Eur. J. Immunol.* 25: 466-473.
4. Bell, M.D. et al. (1994) Upregulation of the macrophage scavenger receptor in response to different forms of injury in the CNS. *J. Neurocytol.* 23: 605-613.
5. Hughes, D.A. et al. (1994) Murine Macrophage Scavenger Receptor: Adhesion function and Expression. *Imm. Letts.* 43: 7-14.
6. Rosen, H. and Hughes, D.A. (1995) Assays of Myeloid Cell Function: Migration and adhesion in vivo. *Weir Handbook of Experimental Immunology*. London, Blackwell Scientific Publications. 5th, ed. In Press.
7. Daugherty, A. et al. (2000) Polymorphism of class A scavenger receptors in C57BL/6 mice. *J. Lipid Res* 41: 1568 - 1577.
8. Moldenhauer, L.M. et al. (2010) GM-CSF Is an Essential Regulator of T Cell Activation Competence in Uterine Dendritic Cells during Early Pregnancy in Mice. *J Immunol.* 185: 7085-7096.
9. Luechtenborg, B. et al. (2008) Function of scavenger receptor class A type I/II is not important for smooth muscle foam cell formation. *Eur J Cell Biol.* 87: 91-9.
10. Sever-Chroneos, Z. et al. (2011) Surfactant Protein A (SP-A)-mediated Clearance of *Staphylococcus aureus* Involves Binding of SP-A to the Staphylococcal Adhesin Eap and the Macrophage Receptors SP-A Receptor 210 and Scavenger Receptor Class A. *J Biol Chem.* 286: 4854-70.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com