

Monoclonal Antibody to Rat IgG1 (heavy chain) - HRP

Catalog No.:	SM1030HRP
Quantity:	0.5 mg
Concentration:	1.0 mg/ml
Background:	Immunoglobulin G (IgG), is one of the most abundant proteins in human serum with normal levels between 8-17 mg/ml in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10e11 variants. Immunoglobulins, or antibodies, are a complex, heterogeneous mixture of proteins that exhibit two fundamental types of structural variation. Subtle structural differences in their antigen combining sites, or variable regions, account for their unique antigen binding specificities.
Host / Isotype:	Mouse / IgG1
Clone:	MARG1.2
Format:	State: Liquid purified IgG fraction. Purification: Affinity Chromatography. Buffer System: PBS, pH 7.4 containing 0.01% Thiomersal as preservative and 50% Glycerol as stabilizer. Label: HRP – Horseradish Peroxidase
Applications:	ELISA: 0.5 µg/ml. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises the gamma-1 immunoglobulin heavy chain and does not bind to other immunoglobulin classes or subclasses. Species: Rat. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General References:	1. Pelegri, C. et al. (2001) Prevention of adjuvant arthritis by the W3/25 anti-CD4 monoclonal antibody is associated with a decrease of blood CD4+ CD45RChigh T cells. Clin. Exp. Immunol. 125: 470-477. 2. Sato, K. et al. (2001) Carbon monoxide generated by heme oxygenase-1 suppresses the rejection of mouse-to-rat cardiac transplants. J. Immunol. 166: 4185-4194.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com