

Monoclonal Antibody to CD4 - PE

Alternate names:	T-cell surface antigen T4/Leu-3, T-cell surface glycoprotein CD4
Catalog No.:	SM1058RT
Quantity:	25 Tests
Concentration:	0.1 mg/ml
Background:	CD4 is a 55kD cell surface glycoprotein that is primarily expressed on a subpopulation of T lymphocytes, on peripheral blood monocytes and on tissue macrophages.
Uniprot ID:	P01730
NCBI:	NP_000607.1
GeneID:	920
Host / Isotype:	Mouse / IgG1
Clone:	RPA-T4
Immunogen:	Human PHA blasts. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Format:	State: Lyophilized purified IgG fraction. Purification: Affinity Chromatography on Protein G Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer. Label: PE – R. Phycoerythrin (RPE) Reconstitution: Restore with distilled water
Applications:	Flow Cytometry: Use 10 µl of neat-1/10 diluted antibody to label 10e6 cells or 100 µl of whole blood. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises CD4. Epitope mapping studies have shown that antibodies, produced by clone RPA-T4, recognise an epitope within domain 1, of the extracellular region, of the CD4 molecule. Clone RPA-T4 has been reported to block gp120-CD4 interaction and inhibit syncytium formation. We recommend the use of SM1058LE for this purpose. Species: Human. Other species not tested.
Storage:	Prior to and following reconstitution store the antibody at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.


For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

- General References:**
1. Zarkesh-Esfahani, H. et al. (2001) High-dose leptin activates human leukocytes via receptor expression on monocytes. *J. Immunol.*167:4593 - 4599.
 2. Voehringer, D. et al. (2002) Lack of proliferative capacity of human effector and memory T cells expressing killer cell lectin-like receptor G1 (KLRG1). *Blood.*100: 3698 - 3702.
 3. Piatier-Tonneau, D. (1997) CD4 workshop panel report. In *Leucocyte Typing VI. White cell differentiation antigens*. Edited by Kishimoto, T., Kikutani, H., von dem Borne, A.E.G.Kr., Goyert, A.M., Mason, D.Y., Miyasaka, M., Moretta, L., Okumura, K., Shaw, S., Springer, T.A., Sugamura, K., Zola, H. Garland publishing Inc. New York & London.

Pictures:



Staining of human peripheral blood lymphocytes with MOUSE ANTI HUMAN CD4:RPE (SM1058R).

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com