

Monoclonal Antibody to CD38 - FITC

Alternate names: ADP-ribosyl cyclase 1, Cyclic ADP-ribose hydrolase 1, T10, cADPr hydrolase 1

Catalog No.: SM1106F
Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: CD38 is expressed by plasma cells, monocytes, early lymphoid cells and activated T cells.

CD38 is widely used to study the processes of T and B cell differentiation and activation.

Recently CD38 has been shown to possess ADP-ribosyl cyclase activity.

Uniprot ID: P28907

NCBI: <u>NP_001766.2</u>

GenelD: 952

Host / Isotype: Mouse / IgG1 Clone: AT13/5

Immunogen: Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1

myeloma cell line.

Format: State: Liquid purified IgG fraction.

Purification: Affinity Chromatography on Protein A.

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as

stabilizer.

Label: FITC - Fluorescein Isothiocyanate Isomer 1

Applications: Flow Cytometry.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises the CD38 cell surface antigen, a 45kD glycoprotein.

Species: Human.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General References: 1. Malavasi, F. et al. (1994) Human CD38: a glycoprotein in search of a function. Immunol.

Today. 15: 95-97.


2. Ellis, J.H. et al. (1995) Engineered anti-CD38 monoclonal antibodies for immunotherapy

of multiple myeloma. J. Immunol. 155: 925-937.





Pictures:



Staining of human peripheral blood monocytes using Mouse Anti Human CD38-FITC