

Monoclonal Antibody to CD45 / LCA (CD45RO) - Purified

Alternate names:	L-CA, Leukocyte common antigen, PTPRC, Receptor-type tyrosine-protein phosphatase C, T200
Catalog No.:	SM1117PS
Quantity:	0.1 mg
Concentration:	1.0 mg/ml
Background:	The antigen is expressed by a functional subset of T cells with memory phenotype. In peripheral blood UCHL1 stains 40-80% of lymphocytes, and all monocytes and granulocytes.
Uniprot ID:	P08575
NCBI:	NP_002829.2
GeneID:	5788
Host / Isotype:	Mouse / IgG2a
Clone:	UCHL1
Immunogen:	Human IL-2 dependent T-Cells. Remarks: Spleen cells from immunised mice were fused with cells of the mouse P3/NS1/1-Ag4-1 myeloma cell line.
Format:	State: Liquid purified IgG fraction. Purification: Affinity Chromatography on Protein G. Buffer System: PBS containing 0.09% Sodium Azide as preservative.
Applications:	Flow Cytometry: Use 10 µl of 1/25-1/50 diluted antibody to label 10e6 cells in 100 µl. Immunoprecipitation. Immunohistochemistry on Frozen Sections. Immunohistochemistry on Paraffin Sections: This product does not require protein digestion pre-treatment of paraffin sections and does not require antigen retrieval using heat treatment prior to staining of paraffin sections. <i>Recommended Positive Control Tissue:</i> Tonsil. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises the low molecular weight isoform (180kD) of the Leucocyte Common Antigen (LCA). Species: Human and Canine. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch


For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

- General References:**
1. Smith, S.H. et al. (1986) Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. *Immunology* 58: 63-70.
 2. Norton, A.J. et al. (1986) Monoclonal antibody (UCHL1) that recognises normal and neoplastic T-cells in routinely fixed tissues. *J. Clin. Pathol.* 39: 399-405.
 3. Beverley, P.C.L. et al. (1986) T-cell subsets and function. *Progress in Immunology VI*. Cinader, B., Miller, G.G., eds. Academic Press Orlando pp 941-948.
 4. Beverley, P.C.L. (1987) Human Tcell subsets. *Immunology Letters* 14: 263-267.
 5. Terry, L.A. et al. (1987) Phenotypic heterogeneity of the CD4+ and CD8+ subsets. *Leucocyte Typing III*. McMichael, A.J., Beverley, P.C.L. et al. eds. University Press. pp 225-227.
 6. Akbar, A.N. et al. (1988) Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T-cells. *J. Immunol.* 140: 2171 - 2178
 7. Terry, L.A. et al. (1988) The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45 *Immunology* 64: 331-336.
 8. Beverley, P.C.L. et al. (1988) Phenotypic diversity of the CD45 antigen and its relation to function. *Immunology, Suppl.* 1: 3-5.
 9. Merckenschlager, M. et al. (1988) Limited dilution analysis of proliferative responses in human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T-cell memory formation. *Eur. J. Immunol.* 18: 1653-1661.
 10. Cavers, M. et al. (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. *Clin Exp Immunol.* 127: 60-5.
 11. Hutnick, N.A. et al. (2010) Vaccination with Ad5 vectors expands Ad5-specific CD8 T cells without altering memory phenotype or functionality. *PLoS One.* 5: e14385.
 12. Leigh, J.E. et al. (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. *Clin Vaccine Immunol.* 13: 678-83.

Pictures:



Staining of human peripheral blood lymphocytes with Mouse Anti Human CD45RO (SM1117PS).

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com