

Monoclonal Antibody to CD62L / L-Selectin - FITC

Alternate names: CD62 antigen-like family member L, LAM-1, LECAM1, LNHR, LYAM1, Leu-8, Leukocyte

adhesion molecule 1, Leukocyte surface antigen Leu-8, Leukocyte-endothelial cell

adhesion molecule 1, Lymph node homing receptor, SELL, TQ1, gp90-MEL

Catalog No.: SM1149F
Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: CD62L is a 74-95kDa member of the selectin family of adhesion receptors, which acts as a

ligand for both CD62-P (P-selectin) and CD62E (E-selectin). Human CD62L is constitutively expressed on most leucocytes including monocytes, granulocytes, lymphocytes, NK cells, bone marrow myeloid progenitor cells and on a subset of thymocytes. CD62L plays an important role in leucocyte tethering and rolling on the endothelial cell surface and for the homing of naive lymphocytes to lymph nodes and Peyers patches via HEV. Neutrophils require a constant supply of this molecule on the cell surface for migration into peripheral tissues and adhesion to activated endothelium at sites of inflammation, where CD62L is rapidly shed as soluble L-selectin, but surface expression still remains. The expression of

CD62L is down regulated on lymphocytes and neutrophils by PMA stimulation.

Uniprot ID: P14151

NCBI: NP 000646.2

GenelD: <u>6402</u>

Host / Isotype: Mouse / IgG2b

Clone: FMC46

Immunogen: PHA stimulated lymphoblasts. Spleen cells from immunised BALB/c mice were fused with

cells of the mouse NS1 myeloma cell line.

Format: State: Liquid purified IgG fraction.

Purification: Affinity Chromatography on Protein G.

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as

stabilizer.

Label: FITC - Fluorescein Isothiocyanate Isomer 1

Applications: Flow Cytometry.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognizes CD62L, also known a L-selectin.

Species: Human, Cynomolgus Monkey, Bovine and Rhesus Monkey.

Other species not tested.



SM1149F: Monoclonal Antibody to CD62L / L-Selectin - FITC

Storage:


Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General References: 1. Pilarski, L. M. et al. (1991) FMC46, a cell protrusion-associated leucocyte adhesion molecule-1 epitope on human lymphocytes and thymocytes. J. Immunol. 147: 136-143. 2. Zola, H. et al. (1991) The expression of sub-population markers on B cells: A re-evaluation using high sensitivity fluorescence flow cytometry. Dis. Markers 9: 103-118. 3. Sopp, P. and Howard, C. C. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. Vet. Immunol. Immunopathol. 56: 11-25.

> 4. Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (Macaca fascicularis) by using anti-human cross-reactive antibodies. Exp. Anim. 49: 97-110.

Pictures:



Staining of human peripheral blood granulocytes with Mouse Anti Human CD62L-FITC (SM1149F/FT).