

Monoclonal Antibody to CD90 - PE

Alternate names:	CDw90, THY1, Thy-1, Thy-1 membrane glycoprotein
Catalog No.:	SM1170RT
Quantity:	25 Tests
Background:	CD90 antigen is a 25 kD glycoprotein homologous to rat Thy-1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by some cells within the thymus. CD90 is also expressed extensively within the brain.
Uniprot ID:	P04216
NCBI:	NP_006279.2
GeneID:	7070
Host / Isotype:	Mouse / IgG1
Clone:	F15-42-1
Immunogen:	Purified Human brain Thy-1. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Format:	State: Lyophilized purified IgG fraction. Purification: Ion Exchange Chromatography. Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA, 5% Sucrose as stabilizer. Label: PE – R. Phycoerythrin (RPE) Reconstitution: Restore in 0.25 ml distilled water. Note. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. We recommend that the vial is gently vortexed after reconstitution and microcentrifuged before use.
Applications:	Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells in 100 µl. This product is routinely tested in Flow Cytometry on the MOLT4 cell line. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	SM1170R recognises the Human CD90 cell surface antigen. Species: Human. Reacts with Cynomolgus Monkey. Other species not tested.
Storage:	Prior to and following reconstitution store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.


For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

- General References:**
1. McKenzie, J.L. and Fabre, J.W. (1981) Human Thy-1: Unusual localization and possible functional significance in lymphoid tissues. *J. Immunol.* 126: 843-850.
 2. Daar, A.S. and Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in human breast carcinomas. *Lancet.* 2: 434-438.
 3. Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. *Exp. Anim.* 49 (2): 97-110.
 4. Fiegat, H. C. et al. (2004) Stem-like cells in human hepatoblastoma. *J. Histochem. Cytochem.* 52: 1495-1501.
 5. Hagood, J. S. et al. (2005) Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. *Am. J. Pathol.* 167: 365-379.
 6. Diaz-Romero, J. et al. (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. *J Cell Physiol.* 214: 75-83.

Pictures:



Staining of HUT78 cells with anti Human CD90-RPE (SM1170RT).

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com