

Monoclonal Antibody to CD89 / FCAR - PE

Alternate names: IgA Fc receptor, Immunoglobulin alpha Fc receptor

Catalog No.: SM1198RT Quantity: 25 Tests

Background: CD89 cell surface antigen is a 50-70kD glycoprotein that is also known as the IgA receptor.

CD89 is expressed by peripheral blood neutrophils and monocytes.

Uniprot ID: P24071

NCBI: NP_001991.1

GenelD: <u>2204</u>

Host / Isotype: Mouse / IgG1 Clone: MIP8a

Immunogen: Recombinant soluble Human Fc alpha R.

Format: State: Lyophilized purified IgG fraction.

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as

stabilizer.

Label: PE – R. Phycoerythrin (RPE)

Reconstitution: Restore with 1 ml distilled water

Applications: Flow Cytometry: Use 10 μl of neat CD89 antibody to label 10e6 cells or 100 μl whole blood.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognizes the Human CD89 cell surface antigen.

Species: Human.

Other species not tested.

Storage: Store the antibody undiluted (Prior to and following reconstitution) at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.

General References: 1. Zhang, W. et al. (2000) Neutrophil lactoferrin release induced by IgA immune complexes

differed from that induced by cross-linking of Fc alpha receptors (Fc alpha R) with a

monoclonal antibody, MIP8a. Clin. Exp. Immunol. 121: 106-111.


2. Lu, J. et al. (2011) Recognition and functional activation of the human IgA receptor

(Fc{alpha}RI) by C-reactive protein. Proc Natl Acad Sci U S A. 108: 4974-9.





Pictures:



Staining of peripheral blood granulocytes with Mouse anti Human CD89 antibody RPE conjugated (SM1198R).