

Monoclonal Antibody to CD95 / FAS - FITC

Alternate names:	APT1, Apo-1 antigen, FAS1, FASLG receptor, TNFRSF6, Tumor necrosis factor receptor superfamily member 6
Catalog No.:	SM1203FT
Quantity:	25 µg
Concentration:	0.1 mg/ml
Background:	<p>CD95, also known as FAS or APO1, is a 36 kDa cell surface type I membrane glycoprotein with an apparent molecular weight of 44 kDa on SDS PAGE. CD95 is a member of the TNF receptor family, which includes TNFR1, TNFR2, CD27, CD30 and CD40. Binding of CD95 Ligand to CD95 or crosslinking of CD95 by anti CD95 monoclonal antibodies leads to apoptosis of CD95 expressing cells. CD95 belongs to a subgroup of family members that have a death domain (DD) which contains 70 amino acids near the carboxyl terminal region of the molecule. The binding of adaptor molecules to this DD is responsible for transmitting the death signal for apoptosis. Stimulation of CD95 results in aggregation of its DD, leading to the recruitment of FADD and caspase 8 that together with the receptor form the death inducing signaling complex (DISC). CD95/CD95L is involved in the peripheral deletion of activated mature T cells at the end of the immune response and defects in this pathway predispose to autoimmune disorders. CD95 is also involved in killing of targets such as virus infected cells or cancer cells and killing of inflammatory cells at immune privileged sites.</p> <p>CD95 is expressed by activated T and B cells, NK cells and thymocytes.</p>
Uniprot ID:	P25445
NCBI:	NP_000034.1
GeneID:	355
Host / Isotype:	Mouse / IgG1
Clone:	LOB 3/17
Immunogen:	Fusion protein comprising extracellular domain of human Fas linked to human Fc. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Format:	<p>State: Liquid purified IgG fraction.</p> <p>Purification: Affinity Chromatography on Protein A.</p> <p>Buffer System: PBS, pH7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer.</p> <p>Label: FITC – Fluorescein Isothiocyanate Isomer 1</p>
Applications:	Flow Cytometry: Use 10 µl of neat-1/10 diluted antibody to label 10e6 cells in 100 µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.

For research and in vitro use only. Not for diagnostic or therapeutic work.


Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Specificity: This antibody recognises the CD95 cell surface antigen.
Species: Human and Rhesus Monkey.
Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.
This product is photosensitive and should be protected from light.
Avoid repeated freezing and thawing.
Shelf life: one year from despatch.

Pictures:



SM1203F/FT (Mouse anti Human CD95 Antibody -FITC) staining of HUT78 T cell line.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com