

Monoclonal Antibody to CD19 - Biotin

Alternate names: B-cell marker, B-lymphocyte surface antigen B4, Differentiation antigen CD19, Leu-12

Catalog No.: SM1529B

Quantity: 0.1 mg

Concentration: 0.1 mg/ml

Background: Human CD19 cell surface antigen is a 96 kD single chain glycoprotein on the surface of

human B lineage cells, including pre-B cells. B cells can be detected in peripheral blood

and tissues.

Uniprot ID: P15391

NCBI: <u>NP_001171569.1</u>

GenelD: <u>930</u>

Host / Isotype: Mouse / IgG1

Clone: LT19

Format: State: Liquid purified IgG fraction.

Purification: Affinity chromatography on Protein G.

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as

stabilizer. **Label:** Biotin

Applications: Flow Cytometry.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises the CD19 cell surface antigen. It labels all B lymphocytes in

peripheral blood and spleen but is negative on granulocytes, monocytes, platelets, T

lymphocytes and erythrocytes.

The antibody reacts with most B cell malignancies including chronic leukaemia and most

lymphomas, but not multiple myeloma.

Species: Human.


Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.



Pictures:



Staining of human peripheral blood lymphocytes with Mouse Anti Human CD19-Biotin (SM1529B/BT).