

Monoclonal Antibody to CD31 / PECAM1 - PE

Alternate names: EndoCAM, GPIIA', PECAM-1, Platelet endothelial cell adhesion molecule

Catalog No.: SM1530RT
Quantity: 25 Tests

Background: CD31, also known as platelet endothelial cell adhesion molecule 1 (PECAM1), is a type I

integral membrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is constitutively expressed on the surface of endothelial cells, and concentrated at the junction between them. It is also weakly expressed on many peripheral

lymphoid cells and platelets.

CD31 has been used to measure angiogenesis in association with tumor recurrence. Other studies have also indicated that CD31 and CD34 can be used as markers for myeloid progenitor cells and recognize different subsets of myeloid leukemia infiltrates (granular

sarcomas).

CD31 is a 130kD glycoprotein expressed by platelets, endothelial cells, granulocytes and

some lymphocytes.

Uniprot ID: <u>P16284</u> **NCBI:** 9606

Host / Isotype: Mouse / IgG1

Clone: WM59

Format: State: Lyophilized purified IgG fraction

Purification: Affinity chromatography on Protein G

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as

stabilizer

Label: PE – R. Phycoerythrin (RPE)

Reconstitution: Restore with 1 ml distilled water.

Applications: Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells or 100 µl whole blood.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises the CD31 cell surface antigen.

Species: Human, Cynomolgus monkey, Monkey, Rhesus Monkey.

Other species not tested.



SM1530RT: Monoclonal Antibody to CD31 / PECAM1 - PE

Storage:

Prior to and following reconstitution store the antibody undiluted at 2-8°C.

DO NOT FREEZE!


This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.

General References: 1. Carbanas. C. et al. (1989) Characterisation of a novel myeloid antigen regulated during differentiation of monocytic cells. Eur J. Immunol. 19: 1373-1378.

> 2. Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (Macaca fascicularis) by using anti-human cross-reactive antibodies. Exp. Anim. 49 (2): 97-110

Pictures:



Staining of human peripheral blood platelets with Mouse Anti Human CD31 antibody -RPE (SM1530R/RT).