

Monoclonal Antibody to CD284 / TLR4 - PE

Alternate names:	Toll-like receptor 4
Catalog No.:	SM1677R
Quantity:	100 Tests
Background:	TLR4, also known as CD284, has been demonstrated to act as a receptor for LPS on human monocytes and macrophages. TLR4 signalling of LPS stimulation requires the presence of the MD-2 molecule. TLR4 is weakly expressed by resting cells, but is upregulated following stimulation with LPS.
Uniprot ID:	O00206
NCBI:	NM_138554.3
Host / Isotype:	Mouse / IgG2a
Clone:	HTA125
Immunogen:	Ba/F3 cell line expressing TLR4. Spleen cells from immunised Balb/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Format:	State: Lyophilized purified IgG fraction. Purification: Affinity Chromatography on Protein G. Buffer Solution: PBS with 5% Sucrose Preservatives: 0.09% Sodium Azide Stabilizers: 1% BSA Label: PE – R. Phycoerythrin (RPE) Reconstitution: Restore with 1.0 ml distilled water.
Applications:	Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells or 100 µl whole blood. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises the Toll like receptor 4 (TLR4) cell surface antigen. This antibody has been demonstrated to block activation of monocytes with LPS. We recommend the use of SM1677LE for this purpose.
Species Reactivity:	Tested: Human, Rhesus monkey, Guinea Pig, Porcine and Canine.
Storage:	Prior to and following reconstitution store the antibody at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General References:	1. Shimazu, R. et al. (1999) MD-2, a molecule that confers lipopolysaccharide responsiveness on Toll like receptor 4. J. Exp. Med. 189: 1777-1182. 2. Jiang, Q. et al. (2000) Lipopolysaccharide induces physical proximity between CD14 and

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Toll like receptor 4 (TLR4) prior to nuclear translocation of NF-Kappa B. *J. Immunol.* 165: 3541-3544.


3. Yang, S. et al. (2001) Synergistic effect of muramyl dipeptide with lipopolysaccharide or lipoteichoic acid to induce inflammatory cytokines in human monocytic cells in culture. *Infect. Immun.* 69: 2045-2053.

4. Triantafyllou, M. et al. (2002) Mediators of innate immune recognition of bacteria concentrate in lipid rafts and facilitate lipopolysaccharide-induced cell activation. *J. Cell Sci.* 115: 2603-2611.

5. Kawahara, T. et al. (2001) Type I Helicobacter pylori lipopolysaccharide stimulates toll-like receptor 4 and activates mitogen oxidase 1 in gastric pit cells. *Infect. Immun.* 69: 4382-4839.

6. Devaney, J.M. (2003) Neutrophil elastase up-regulates interleukin-8 via toll-like receptor 4. *FEBS Lett.* 544:129-32.

7. de Kleer, I. (2010) CD30 Discriminates Heat Shock Protein 60-Induced FOXP3+CD4+ T Cells with a Regulatory Phenotype. *J Immunol.* 185(4):2071-9.

Pictures:

Staining of U937 cells with RPE conjugated Mouse Anti Human CD284 antibody (SM1677R).

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com