

US office: Acris Antibodies, Inc. San Diego, CA UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com SM1744P Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

	Monoclonal Antibody to CD45 / LCA - Purified
Alternate names:	L-CA, Leukocyte common antigen, PTPRC, Receptor-type tyrosine-protein phosphatase C, T200
Catalog No.:	SM1744P
Quantity:	0.2 mg
Concentration:	1.0 mg/ml
Background:	CD45 is a family of single chain transmembraneous glycoproteins consisting of at least for isoforms (220, 205, 190, 180 kDa) which share a common large intracellular domain. Their extracellular domains are heavily glycosylated. The different isoforms are produced by alternative messenger RNA splicing of three exons of a single gene on chromosome 1. CD- is expressed on cells of the human hematopoietic lineage (including hematopoietic stem cells) with the exception of mature red cells. It is not detected on differentiated cells of other tissues. It is likely that CD45 plays an important role in signal transduction, inhibitio or upregulation of various immunological functions. Antibodies recognising a common epitope on all of the isoforms are termed CD45RO etc.
Uniprot ID:	<u>P08575</u>
NCBI:	<u>NP_002829.2</u>
GenelD:	<u>5788</u>
Host / Isotype:	Rat / IgG2b
Clone:	YTH24.5
Immunogen:	Human T lymphocytes. Spleen cells from an immunised DA rat were fused with cells of the rat Y3/Ag1.2.3 myeloma cell line.
Format:	State: Liquid purified IgG Purification: Affinity chromatography on Protein A Buffer System: PBS containing 0.09% Sodium Azide
Applications:	Flow Cytometry: Use 10 μ l of 1/10-1/100 diluted antibody to label 10e6 cells in 100 μ l. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises the leukocyte common antigen CD45, expressed on all cells of haematopoietic origin except for erythrocytes. Species: Human. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

OG/20121030

SM1744P: Monoclonal Antibody to CD45 / LCA - Purified

General References: 1. Bindon, C.I. et al. (1985) Therapeutic potential of monoclonal antibodies to the leukocyte common antigen: synergy and interference in complement - mediated lysis. Transplantation 40: 538-544.