

## Monoclonal Antibody to CD44 - PE

<b>Alternate names:</b>	CDw44, ECMR-III, Epican, Extracellular matrix receptor III, GP90 lymphocyte homing/adhesion receptor, HUTCH-I, Heparan sulfate proteoglycan, Hermes antigen, Hyaluronate receptor, LHR, MDU2, MDU3, MIC4, PGP-1, Phagocytic glycoprotein 1
<b>Catalog No.:</b>	SM1898RT
<b>Quantity:</b>	25 Tests
<b>Concentration:</b>	0.1 mg/ml
<b>Background:</b>	CD44 is a 100kD glycoprotein widely expressed on human leucocytes, white matter of the brain and by some epithelial cells of the intestine and of the breast. CD44 expression may be up-regulated upon some carcinomas, and it has been speculated that this may be related to metastatic potential.
<b>Uniprot ID:</b>	<a href="#">P16070</a>
<b>NCBI:</b>	<a href="#">NP_000601.3</a>
<b>GeneID:</b>	<a href="#">960</a>
<b>Host / Isotype:</b>	Mouse / IgG2a
<b>Clone:</b>	F10-44-2
<b>Immunogen:</b>	Human T lymphocytes. Spleen cells of immunised BALB/c mice were fused with cells from the mouse NS1 myeloma line.
<b>Format:</b>	<b>State:</b> Lyophilized purified IgG fraction <b>Purification:</b> Affinity Chromatography on Protein G <b>Buffer System:</b> PBS, pH 7.4, containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer. <b>Label:</b> PE – R. Phycoerythrin (RPE) <b>Reconstitution:</b> Restore with 1 ml distilled water.
<b>Applications:</b>	Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells in 100 µl. This product is routinely tested in flow cytometry on human peripheral blood lymphocytes. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognises the CD44 cell surface antigen. <b>Species:</b> Human, Cynomolgus monkey and Monkey. Other species not tested.
<b>Storage:</b>	Prior to and following reconstitution store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b> This product is photosensitive and should be protected from light. Shelf life: one year from despatch.


**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)

- General References:**
1. Dalchau, R. et al. (1980) Monoclonal antibody to a human brain granulocyte T lymphocyte antigen probably homologous to the W3/13 antigen of the rat. *Eur. J. Immunol.* 10: 745 - 750.
  2. Daar, A.S. and Fabre, J.W. (1981) Demonstration of an unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in human breast carcinoma. *Lancet.* 2: 434-438.
  3. Cattoretti, G. et al. (1993) Antigen unmasking on formalin-fixed, paraffin-embedded tissue sections. *J. Pathol.* 171:83-98.
  4. Yoshino, N. et al. (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of Cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. *Exp. Anim.* 49 (2): 97-110.
  5. Roscic - Mrkic, B. et al. (2003) RANTES (CCL5) uses the proteoglycan CD44 as an auxiliary receptor to mediate cellular activation signals and HIV - 1 enhancement. *Blood.* 102: 1169 - 1177.

**Pictures:**



Staining of human peripheral blood lymphocytes with MOUSE ANTI HUMAN CD44:RPE (SM1898R).

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)