

Monoclonal Antibody to CD31 / PECAM1 - FITC

Alternate names:	EndoCAM, GPIIA', PECAM-1, Platelet endothelial cell adhesion molecule
Catalog No.:	SM272F
Quantity:	0.1 mg
Concentration:	0.1 mg/ml
Background:	<p>CD31, also known as platelet endothelial cell adhesion molecule 1 (PECAM1), is a type I integral membrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is constitutively expressed on the surface of endothelial cells, and concentrated at the junction between them. It is also weakly expressed on many peripheral lymphoid cells and platelets.</p> <p>CD31 has been used to measure angiogenesis in association with tumor recurrence. Other studies have also indicated that CD31 and CD34 can be used as markers for myeloid progenitor cells and recognize different subsets of myeloid leukemia infiltrates (granular sarcomas).</p>
Uniprot ID:	Q3SWT0
NCBI:	NP_113779.1
GeneID:	29583
Host / Isotype:	Mouse / IgG1
Clone:	TLD-3A12
Immunogen:	Activated, Lewis rat derived microglial cells. Spleen cells from immunised BALB/c mouse were fused with cells of the mouse SP2 myeloma cell line.
Format:	State: Liquid purified IgG fraction Purification: Affinity Chromatography on Protein G. Buffer System: PBS, pH 7.4 containing 1% BSA as stabilizer and 0.09% Sodium Azide as preservative Label: FITC – Fluorescein Isothiocyanate Isomer I
Applications:	Flow Cytometry: Use 10 µl of neat antibody to label 10e6 cells in 100 µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises PECAM-1 (CD31), a type 1 transmembrane protein expressed primarily on endothelial cells, platelets and leucocytes. Clone TLD-3A12 has been shown to partially block the proliferative response of antigen-specific CD4+ T cells to antigen-presenting cells and relevant antigen. (3) We recommend SM272LE for use in functional studies. Species: Rat. Other species not tested.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Storage:

Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.
This product is photosensitive and should be protected from light.
Avoid repeated freezing and thawing.
Shelf life: one year from despatch.

General References:

1. Flaris, N.A., Densmore, T.L., Molleston, M.C. and Hickey, W.F. (1993). Characterisation of microglia and macrophages in the central nervous system of the rat. *GLIA* 7: 34-40.
2. Male, D., Rahman, J., Linke, A., Zhao, W. and Hickey, W.F. (1995). An interferon inducible molecule on brain endothelium which controls lymphocyte adhesion mediated by integrins. *Immunology* 84: 453-460.
3. Williams, K.C. et al. (1996) PECAM-1 (CD31) expression in the central nervous system and its role in experimental allergic encephalomyelitis in the rat. *J. Neurosci. Res.* 45(6):747-57.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com