

## Monoclonal Antibody to CD172a / SIRPA - FITC

<b>Alternate names:</b>	BIT, Brain Ig-like molecule with tyrosine-based activation motifs, CD172 antigen-like family member A, Inhibitory receptor SHPS-1, MFR, MYD1, Macrophage fusion receptor, MyD-1 antigen, PTPNS1, SHP substrate 1, SHPS-1, SHPS1, Signal-regulatory protein alpha-1, Sirp-alpha-2, Sirp-alpha-3, Tyrosine-protein phosphatase non-receptor type substrate 1, p84
<b>Catalog No.:</b>	SM275F
<b>Quantity:</b>	0.1 mg
<b>Concentration:</b>	0.1 mg/ml
<b>Background:</b>	CD172a is expressed selectively by myeloid cells and by neurons.
<b>Uniprot ID:</b>	<a href="#">P97710</a>
<b>NCBI:</b>	<a href="#">NP_037148.2</a>
<b>GeneID:</b>	<a href="#">25528</a>
<b>Host / Isotype:</b>	Mouse / IgG2a
<b>Clone:</b>	OX-41
<b>Immunogen:</b>	Rat peritoneal macrophages. Spleen cells from immunised BALB/c mice were fused with the NSO/U mouse myeloma cell line.
<b>Format:</b>	<b>State:</b> Liquid purified IgG <b>Purification:</b> Affinity chromatography on Protein G <b>Buffer System:</b> PBS, pH7.4 containing 0.09% Sodium Azide and 1.0% Bovine Serum Albumin <b>Label:</b> FITC – Fluorescein Isothiocyanate Isomer 1
<b>Applications:</b>	Flow Cytometry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognises rat CD172a, also known as Signal regulatory protein (SIRP). MRC OX-41 binds to a different epitope on SIRP than ED9. <b>Species:</b> Rat. Other species not tested.
<b>Storage:</b>	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
<b>General References:</b>	1. Robinson, A.P. et al. (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. Immunology 57: 239-247.

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)

2. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. *Immunology* 42: 593-600.
3. Adams, S. et al. (1998) Signal-regulatory protein is selectively expressed by myeloid and neuronal cells. *J. Immunol.* 161: 1853-1859.

---

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)