

Monoclonal Antibody to CD105 / Endoglin - FITC -


Alternate names:	END, ENG, HHT1, ORW, ORW1
Catalog No.:	SM3078F
Quantity:	100 Tests
Background:	CD105 (Endoglin) is a homodimeric transmembrane glycoprotein serving in presence of TGF β R-2 as a receptor for TGF β -1 and TGF β -3. CD105 is highly expressed on endothelial cells and promotes angiogenesis during wound healing, infarcts and in a wide range of tumours and its gene expression is stimulated by hypoxia. CD105 prevents apoptosis in hypoxic endothelial cells and also antagonises the inhibitory effects of TGF β -1 on vascular endothelial cell growth and migration. Normal cellular levels of CD105 are required for formation of new blood vessels.
Uniprot ID:	P17813
NCBI:	9606
Host / Isotype:	Mouse / IgG2a
Clone:	MEM-226
Immunogen:	Recombinant Vaccinia virus containing the human CD105 cDNA
Format:	State: Liquid purified Ig fraction Buffer System: Phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent. Label: FITC – Conjugated with Fluorescein isothiocyanate
Applications:	Flow Cytometry analysis of human blood cells using 20 μ l reagent / 100 μ l of whole blood or 10e6 cells in a suspension. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	The antibody MEM-226 reacts with CD105 (Endoglin), a 180 kDa type I homodimerizing membrane glycoprotein expressed on vascular endothelial cells (small and large vessels), activated monocytes and tissue macrophages, stromal cells of certain tissues including bone marrow, pre-B lymphocytes in fetal marrow and erythroid precursors in fetal and adult bone marrow; it is also present on syncytiotrophoblast on placenta throughout pregnancy. Species: Human. Other species not tested.
Storage:	Store the antibody at 2 - 8 $^{\circ}$ C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General References:	1. Zhu Y, Sun Y, Xie L, Jin K, Sheibani N, Greenberg DA: Hypoxic induction of endoglin via mitogen-activated protein kinases in mouse brain microvascular endothelial cells. Stroke. 2003 Oct;34(10):2483-8.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

2. Li C, Issa R, Kumar P, Hampson IN, Lopez-Novoa JM, Bernabeu C, Kumar S: CD105 prevents apoptosis in hypoxic endothelial cells. *J Cell Sci.* 2003 Jul 1;116(Pt 13):2677-85.
3. Guo B, Slevin M, Li C, Parameshwar S, Liu D, Kumar P, Bernabeu C, Kumar S: CD105 inhibits transforming growth factor-beta-Smad3 signalling. *Anticancer Res.* 2004 May-Jun;24(3a):1337-45.
4. Warrington K, Hillarby MC, Li C, Letarte M, Kumar S: Functional role of CD105 in TGF-beta1 signalling in murine and human endothelial cells. *Anticancer Res.* 2005 May-Jun;25(3B):1851-64.
5. Piao M, Tokunaga O: Significant expression of endoglin (CD105), TGFbeta-1 and TGFbeta R-2 in the atherosclerotic aorta: an immunohistological study. *J Atheroscler Thromb.* 2006 Apr;13(2):82-9.
6. Schmidt D, Achermann J, Odermatt B, Breyman C, Mol A, Genoni M, Zund G, Hoerstrup SP: Prenatally fabricated autologous human living heart valves based on amniotic fluid derived progenitor cells as single cell source. *Circulation.* 2007 Sep 11;116(11 Suppl):I64-70.

Pictures:

Surface staining of HUVEC (human umbilical vein endothelial cells) with anti-human CD105 (MEM-226) FITC. Total viable cells were used for analysis.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com