

F(ab')₂ Fragment of Rabbit anti-Rat IgG [H&L]

Catalog No.:	SP1021F
Quantity:	1 mg
Concentration:	1.0 mg/ml
Host:	Rabbit
Format:	State: Liquid F(ab') ₂ fragment of IgG Purification: IgG chromatography followed by Pepsin digestion and a gel filtration Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide Label: FITC – Fluorescein Isothiocyanate Isomer I
Applications:	Flow Cytometry: 1/200 - 1/400; Use 50 µl of the suggested working dilution to label 10e6 cells in 100 µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	F(ab') ₂ fragment reacts to IgG. Species: Rat. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General References:	1. O-Charoenrat, P. et al. (2000) Epidermal growth factor-like ligands differentially up-regulate matrix metalloproteinase 9 in head and neck squamous carcinoma cells. Cancer Res. 60: 1121 - 1128.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com