

Product Information Sheet

Product Name: Rabbit Anti-mouse IgG (H+L), Hilyte FluorTM 750-labeled

Catalog Number: 28164-05-H750

Size: 0.5 mg

Concentration: 1 mg/mL

Degree of Substitution:

(DOS)

Dye:protein molar ratio is specified on the vial.

Fluorescence: Excitation/Emission wavelength = 754 nm/778 nm

Storage buffer: 10 mM phosphate, 150 mM NaCl, pH 7.2, with BSA and 2 mM sodium azide as

preservatives.

Storage: The dye-protein conjugate is stable for 2~3 months at 4°C. For long-term

storage, divide the solution into aliquots and store at -20°C or add an equal volume of glycerol (ACS grade or higher) and store the solution at -20°C without aliquoting. Avoid multiple thaw-freeze cycles. The product is stable for

1 year at -20°C.

Instructions: HiLyte FluorTM 750-IgG conjugate has been optimized in fluorophore/protein

labeling ratio to ensure high fluorescent signal and uncompromised IgG function. Spectrally similar to Cy7 dye, HiLyte FluorTM 750 is the longest-wavelength HiLyte FluorTM dye currently available. Its fluorescence emission maximum at 778 nm is well separated from commonly used far-red fluorophores such as HiLyte FluorTM 647, HiLyte FluorTM 680 or allophycocyanin (APC), facilitating multicolor analysis. With a peak excitation at ~754 nm, conjugates of HiLyte FluorTM 750 dyes are well excited by a xenon-arc lamp or dye-pumped

lasers operating in the 720–750 nm range.

The recommended concentration for most immunofluorescent staining is 1-10 μ g/mL. If the non-specific binding background is high, you may decrease the concentration to minimize the background. You may also centrifuge the

conjugate briefly and use the supernatant only for staining.

For in vitro research use only

AnaSpec, Inc. • 34801 Campus Dr • Fremont, CA 94555 Tel: 1-800-452-5530 • Fax 510-791-9572• service@anaspec.com • www.anaspec.com