

# Mucin 5AC (Gastric Mucin)

Concentrated Monoclonal Antibody  
902-231-071717

**BIOCARE**  
M E D I C A L

**Catalog Number:** ACR 231 A  
**Description:** 0.1 ml, concentrated  
**Dilution:** 1:100-1:200  
**Diluent:** Da Vinci Green

## Intended Use:

For Research Use Only. Not for use in diagnostic procedures.

## Summary and Explanation:

Mucins are high molecular weight glycoproteins with 80% carbohydrate contents and the remaining 20% is constituted by protein core. Gastric mucin M1 antigens are found in columnar mucus cells of surface gastric epithelium and in goblet cells of the fetal and pre-cancerous colon, but not in normal colon. Evidence from the literature suggests that they are associated with peptide core of mucins. Resurgence of gastric mucin reactivity during colon carcinogenesis is suggested to be due to either reexpression of the peptide core of gastric (or fetal colonic) mucins in the adult colon, or due to changes on the glycosylation pattern of mucin which expose the hidden M1 antigens.

## Principle of Procedure:

Antigen detection in tissues and cells is a multi-step immunohistochemical process. The initial step binds the primary antibody to its specific epitope. After labeling the antigen with a primary antibody, a secondary antibody is added to bind to the primary antibody. An enzyme label is then added to bind to the secondary antibody; this detection of the bound antibody is evidenced by a colorimetric reaction.

**Source:** Mouse monoclonal

**Species Reactivity:** Human; others not tested

**Clone:** 45M1

**Isotype:** IgG1/kappa

**Total Protein Concentration:** ~10 mg/ml. Call for lot specific Ig concentration.

**Epitope/Antigen:** MUC5AC

**Cellular Localization:** Cytoplasmic

**Positive Control:** Stomach

## Known Applications:

Immunohistochemistry (formalin-fixed paraffin-embedded tissues)

**Supplied As:** Buffer with protein carrier and preservative

## Storage and Stability:

Store at 2°C to 8°C. Do not use after expiration date printed on vial. If reagents are stored under conditions other than those specified in the package insert, they must be verified by the user. Diluted reagents should be used promptly; any remaining reagent should be stored at 2°C to 8°C.

## Protocol Recommendations:

**Peroxide Block:** Block for 5 minutes with Biocare's Peroxidized 1.

**Pretreatment Solution:** Reveal

## Pretreatment Protocol:

Heat Retrieval Method:

Retrieve sections under pressure using Biocare's Decloaking Chamber followed by a wash in distilled water; alternatively, steam tissue sections for 45-60 minutes. Allow solution to cool for 10 minutes then wash in distilled water.

**Protein Block (Optional):** Incubate for 5-10 minutes at RT with Biocare's Background Punisher.

**Primary Antibody:** Incubate for 30 minutes at RT.

**Probe:** Incubate for 10 minutes at RT with a secondary probe.

**Polymer:** Incubate for 10 minutes at RT with a tertiary polymer.

## Protocol Recommendations Cont'd:

### Chromogen:

Incubate for 5 minutes at RT with Biocare's DAB - OR - Incubate for 5-7 minutes at RT with Biocare's Warp Red.

### Counterstain:

Counterstain with hematoxylin. Rinse with deionized water. Apply Tacha's Bluing Solution for 1 minute. Rinse with deionized water.

### Technical Note:

This antibody has been standardized with Biocare's MACH 4 detection system. It can also be used on an automated staining system and with other Biocare polymer detection kits. Use TBS buffer for washing steps.

### Limitations:

This product is provided for Research Use Only (RUO) and is not for use in diagnostic procedures. Suitability for specific applications may vary and it is the responsibility of the end user to determine the appropriate application for its use.

### Precautions:

1. This antibody contains less than 0.1% sodium azide. Concentrations less than 0.1% are not reportable hazardous materials according to U.S. 29 CFR 1910.1200, OSHA Hazard communication and EC Directive 91/155/EC. Sodium azide (NaN<sub>3</sub>) used as a preservative is toxic if ingested. Sodium azide may react with lead and copper plumbing to form highly explosive metal azides. Upon disposal, flush with large volumes of water to prevent azide build-up in plumbing. (Center for Disease Control, 1976, National Institute of Occupational Safety and Health, 1976) (5)
2. Specimens, before and after fixation, and all materials exposed to them should be handled as if capable of transmitting infection and disposed of with proper precautions. Never pipette reagents by mouth and avoid contacting the skin and mucous membranes with reagents and specimens. If reagents or specimens come in contact with sensitive areas, wash with copious amounts of water. (6)
3. Microbial contamination of reagents may result in an increase in nonspecific staining.
4. Incubation times or temperatures other than those specified may give erroneous results. The user must validate any such change.
5. Do not use reagent after the expiration date printed on the vial.
6. The MSDS is available upon request and is located at <http://biocare.net/support/msds/>.

### Technical Support:

Contact Biocare's Technical Support at 1-800-542-2002 for questions regarding this product.

### References:

1. Fetsch PA, Abati A, Hijazi YM. Utility of the antibodies CA 19-9, HBME-1, and thrombomodulin in the diagnosis of malignant mesothelioma and adenocarcinoma in cytology. *Cancer*. 1998 Apr 25;84 (2):101-8.
2. Ichihara T, *et al*. Immuno-histochemical localization of CA 19-9 and CEA in pancreatic carcinoma and associated diseases. *Cancer*. 1988 Jan 15;61(2):324-33.
3. Soslow RA, *et al*. Transitional cell neoplasms of the ovary and urinary bladder: a comparative immuno- histochemical analysis. *Int J Gynecol Pathol*. 1996 Jul;15(3):257 -65.



60 Berry Drive  
Pacheco, CA 94553  
USA

Rev: 062117

Tel: 800-799-9499 | [www.biocare.net](http://www.biocare.net) | Fax: 925-603-8080

## Mucin 5AC (Gastric Mucin)

Concentrated Monoclonal Antibody  
902-231-071717

**BIOCARE**  
M E D I C A L

### References Cont'd:

4. Charpin C, *et al.* Carcinoembryonic antigen (CEA) and carbohydrate determinant 19 -9 (CA 19-9) localization in 121 primary and metastatic ovarian tumors: an immunohistochemical study with the use of monoclonal antibodies. *Int J Gynecol Pathol.* 1982;1(3):231-45.
5. Center for Disease Control Manual. Guide: Safety Management, NO. CDC-22, Atlanta, GA. April 30, 1976 "Decontamination of Laboratory Sink Drains to Remove Azide Salts."
6. Clinical and Laboratory Standards Institute (CLSI). Protection of Laboratory workers from occupationally Acquired Infections; Approved guideline-Third Edition CLSI document M29-A3 Wayne, PA 2005.



60 Berry Drive  
Pacheco, CA 94553  
USA

Rev: 062117

Tel: 800-799-9499 | [www.biocare.net](http://www.biocare.net) | Fax: 925-603-8080