

## Anti-Human CD41a PE

Catalogue Number : 03511-60

RUO: For Research Use Only. Not for use in diagnostic procedures.

---

### Product Information

**Clone:** HIP8

**Format/Conjugate:** PE

**Concentration:** 5 uL (0.125 ug)/test

**Reactivity:** Human

**Laser:** Blue (488nm), Yellow/Green (532-561nm)

**Peak Emission:** 578nm

**Peak Excitation:** 496nm

**Filter:** 585/40

**Brightness (1=dim,5=brightest):** 5

**Isotype:** Mouse IgG1, kappa

**Formulation:** Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, pH7.2.

**Storage:** Product should be kept at 2-8°C and protected from prolonged exposure to light.

**Applications:** FC

### Description

The HIP8 monoclonal antibody specifically reacts with human CD41 (GPIIb), a subunit of the calcium-dependent complex with CD61 (IIIa). The complex is expressed on megakaryocytes and platelets. The HIP8 antibody blocks the aggregation of platelets and inhibits platelet activation.

### Preparation & Storage

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

### Application Notes

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. The antibody can be used at less than or equal to 5 µL per test. A test is the amount of antibody required to stain a cell sample in the final volume of 100 µL.

### References

1. Leucocyte typing IV: white cell differentiation antigens. Oxford University Press, 1989.
2. Schlossman, S. F. (1995).; Leucocyte typing V: White cell differentiation antigens: Proceedings of the Fifth International Workshop and Conference, Held in Boston, USA 3-7 November, 1993. Oxford University Press.