



# Anti-Human/Mouse CD44 FITC

Catalogue Number: 06511-50

RUO: For Research Use Only. Not for use in diagnostic procedures.

#### **Product Information**

Clone: IM7

Format/Conjugate: FITC Concentration: 0.5 mg/mL Reactivity: Human, Mouse Laser: Blue (488nm) Peak Emission: 520nm Peak Excitation: 494nm

Filter: 530/30

Brightness (1=dim,5=brightest): 3

Isotype: Rat IgG2b, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

Applications: FC

#### Description

The IM7 monoclonal antibody specifically reacts with all the isoforms and both alloantigens of the CD44 glycoprotein (Pgp-1, Ly-24). CD44 is expressed on hematopoietic and non-hematopoietic cells, bone marrow myeloid cells, memory T lymphocytes, periphery activated B cells, CD4+ T lymphocytes, and CD8+ T lymphocytes. The periphery B and T lymphocytes upregulate the expression of CD44. CD44 binds to hyaluronan molecules, acting as an adhesion molecule. The IM7 antibody inhibits collagen-induced arthritis in DBA/1 mice, prevents central nervous system inflammation associated with experimental autoimmune encephalomyelitis, but exacerbates the experimental autoimmune thyroiditis in CRA/1 mice.

The IM7 antibody also cross-reacts with dog, human, pig, horse, cat, and cow leukocytes.

## **Preparation & Storage**

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

### **Application Notes**

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. For flow cytometric staining, the suggested use of this reagent is  $\leq$ 0.5 ug per million cells in 100  $\mu$ l volume. It is recommended that the reagent be titrated for optimal performance for each application.

## References

- 1.Trowbridge, I. S., Lesley, J., Schulte, R., Hyman, R., ; Trotter, J. (1982). Biochemical characterization and cellular distribution of a polymorphic, murine cell-surface glycoprotein expressed on lymphoid tissues.; Immunogenetics, 15(3), 299-312.
- 2. Lesley, J., ; Trowbridge, I. S. (1982). Genetic characterization of a polymorphic murine cell-surface glycoprotein.; Immunogenetics,;15(3), 313-320.
- 3. Nedvetzki, S., Walmsley, M., Alpert, E., Williams, R. O., Feldmann, M., ; Naor, D. (1999). CD44 involvement in experimental collagen-induced arthritis (CIA).; Journal of autoimmunity,; 13(1), 39-47.