



# Anti-Human CD15 FITC

Catalogue Number: 07211-50

RUO: For Research Use Only. Not for use in diagnostic procedures.

#### **Product Information**

Clone: HI98

Format/Conjugate: FITC

Concentration: 5 uL (0.125 ug)/test

Reactivity: Human
Laser: Blue (488nm)
Peak Emission: 520nm
Peak Excitation: 494nm

Filter: 530/30

Brightness (1=dim,5=brightest): 3

Isotype: Mouse IgM, kappa

Formulation: Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.

Storage: Product should be kept at 2-8°C and protected from prolonged exposure to light.

Applications: FC

### Description

The HI98 monoclonal antibody specifically reacts with CD15, a 220 kDa carbohydrate moiety also known as SSEA-1, Lewis 3-fucosyl-N-acetyllactosamine (3-FAL). CD15 is expressed on most granulocytes and monocytes to the exclusion of lymphocytes and basophils. It is involved in cell adhesion, chemotaxis, and phagocytosis.

### **Preparation & Storage**

The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.

## **Application Notes**

The antibody has been analyzed for quality through the flow cytometric analysis of the relevant cell type. The antibody can be used at less than or equal to 5  $\mu$ L per test. A test is the amount of antibody required to stain a cell sample in the final volume of 100  $\mu$ L.

### References

 ${\it 1.} Leucocyte\ typing\ IV:\ white\ cell\ differentiation\ antigens.\ Oxford\ University\ Press,\ 1989.$ 

- 2. Barclay, A. N., Brown, M. H., Law, S. A. K. A., McKnight, A. J., Tomlinson, M. G., ; van der Merwe, P. A. (1997).; The leucocyte antigen factsbook. Academic Press.
- 3. Lund-Johansen, F., Olweus, J., Horejsi, V., Skubitz, K. M., Thompson, J. S., Vilella, R., ; Symington, F. W. (1992). Activation of human phagocytes through carbohydrate antigens (CD15, sialyl-CD15, CDw17, and CDw65).; Journal of immunology (Baltimore, Md.: 1950),;148(10), 3221.