

bs-0432R-Cy5**[Conjugated Secondary Antibody]**

www.biossusa.com

support@biossusa.com

800.501.7654 [DOMESTIC]

+1.781.569.5821 [INTERNATIONAL]

Rabbit Anti-Chicken IgY Antibody, Cy5 Conjugated**DATASHEET****Host:** Rabbit**Target Protein:** Chicken IgY**Clonality:** Polyclonal**Isotype:** IgG**Source:** Chicken IgY**Purification:** Purified by Protein A.**Storage:** Aqueous buffered solution containing 1% BSA, 50% glycerol and 0.09% sodium azide. Store at 4°C for 12 months.

Background: In chickens, immunoglobulin Y is the functional equivalent to Immunoglobulin G (IgG). Like IgG, it is composed of two light and two heavy chains. Structurally, these two types of immunoglobulin differ primarily in the heavy chains, which in IgY have a molecular mass of about 65,100 atomic mass units (amu), and are thus larger than in IgG. The light chains in IgY, with a molar mass of about 18,700 amu, are somewhat smaller than the light chains in IgG. The molar mass of IgY thus amounts to about 167,000 amu. The steric flexibility of the IgY molecule is less than that of IgG. Functionally, IgY is partially comparable to Immunoglobulin E (IgE), as well as to IgG. However, in contrast to IgG, IgY does not bind to Protein A, to Protein G, or to cellular Fc receptors. Furthermore, IgY does not activate the complement system. The name Immunoglobulin Y was suggested in 1969 by G.A. Leslie and L.W. Clem, after they were able to show differences between the immunoglobulins found in chicken eggs, and immunoglobulin G. Other synonymous names are Chicken IgG, Egg Yolk IgG, and 7S-IgG.

Conjugation: Cy5**Excitation/** 625,650nm/670nm**Emission:****Size:** 200ul**Concentration:** 1ug/ul**Applications:** IF(IHC-P)(1:500-2000)**Cross Reactive** Chicken
Species:

For research use only. Not intended for diagnostic or therapeutic use.