



6044 Cornerstone Court W, Ste E
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet

TNKS1 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺) **Catalog # 80579**

DESCRIPTION: The TNKS1 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺) is designed to measure Tankyrase 1 (TNKS1) activity for screening and profiling applications. TNKS1 catalyzes the NAD-dependent addition of poly(ADP-ribose) to the substrate proteins. The TNKS1 assay kit comes in a convenient 384-well format, with purified TNKS1 enzyme, histone mixture, and PARP assay buffer for 384 enzyme reactions. The key to the TNKS1 Histone Ribosylation Assay (Biotin-labeled NAD⁺) is the biotinylated substrate. With this kit, only three simple steps are required for TNKS1 reactions. First, histone proteins are coated on a 384-well plate. Next, the biotinylated TNKS1 substrate is incubated with an assay buffer that contains the TNKS1 enzyme. Finally, the plate is treated with streptavidin-HRP followed by addition of the HRP substrate to produce chemiluminescence that can be measured using a chemiluminescence reader.

COMPONENTS:

| Reagent | Amount | Storage |
|---|------------|------------------|
| TNKS1 (BPS # 80504) | 10 µg | -80 °C |
| 5x histone mixture | 2 ml | -80 °C |
| 10x Assay Mixture containing biotinylated substrate | 600 µl | -80 °C |
| 10x PARP assay buffer | 2 ml | -20 °C |
| Blocking buffer | 50 ml | +4 °C |
| Streptavidin-HRP | 200 µl | +4 °C |
| HRP chemiluminescent substrate (2 components) | 12 ml each | +4 °C |
| 384-well plate | 1 | Room Temperature |

MATERIALS AND INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

1x PBS buffer
PBST buffer (1x PBS, containing 0.05% Tween-20)
Luminometer or fluorescent microplate reader capable of reading chemiluminescence
Adjustable micropipettor and sterile tips
Rotating or rocker platform

APPLICATIONS: Great for studying enzyme kinetics and screening small molecular inhibitors for drug discovery and HTS applications.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6044 Cornerstone Court W, Ste E
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

STABILITY: Up to 1 year when stored as recommended.

REFERENCE:

Brown, J.A., Marala, R.B. *J. Pharmacol. Toxicol. Methods* 2002 **47**:137-41.

Assay Protocol:

All samples and controls should be tested in duplicate.

Coating the plate with the histone mixture:

- 1) Dilute 5x histone mixture 1:5 in PBS.
- 2) Add 25 μ l diluted histone mixture to each well and incubate overnight at 4 °C.
- 3) Wash the plate 3 times with 100 μ l PBST buffer.
- 4) Block the wells by adding 100 μ l of Blocking buffer to every well. Incubate for 60 minutes at room temperature.
- 5) Wash the plate 3 times with 100 μ l PBST buffer

(Alternatively, the plate can be coated for 90 minutes at 37°C followed by 60 minutes blocking at room temperature. All washing steps should be the same.)

Step 1: Ribosylation reaction

- 1) Prepare the master mixture: N wells x (1.25 μ l **10x PARP assay buffer** + 1.25 μ l **10x PARP assay mixture** + 10 μ l **H₂O**)
- 2) Thaw TNKS1 **enzyme** on ice. Upon first thaw, briefly spin tube containing enzyme to recover full content of the tube. Aliquot TNKS1 **enzyme** into single use aliquots. Store remaining undiluted enzyme in aliquots at -80°C immediately. *Note: TNKS1 enzyme is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted enzyme.*
- 3) Dilute TNKS1 **enzyme** in 1X PARP assay buffer at 1.25 – 2.5 ng/ μ l (12.5 - 25 ng/10 μ l). Keep diluted enzyme on ice until use. Discard any unused diluted enzyme after use.

Add 12.5 μ l of master mixture to each well designated for the “Positive Control”, “Test Inhibitor”, and “Blank”. For the “Substrate Control”, add 1.25 μ l **10x PARP assay buffer** + 11.25 μ l **H₂O**.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com



6044 Cornerstone Court W, Ste E
San Diego, CA 92121
Tel: 1.858.829.3082
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

| | Blank | Positive Control | Substrate Control | Test Inhibitor |
|---------------------------------|---------|------------------|-------------------|----------------|
| 10X PARP Assay Buffer | 1.25 µl | 1.25 µl | 1.25 µl | 1.25 µl |
| 10X assay mixture | 1.25 µl | 1.25 µl | – | 1.25 µl |
| H ₂ O | 10 µl | 10 µl | 11.25 µl | 10 µl |
| Test Inhibitor | – | – | – | 2.5 µl |
| Inhibitor buffer (no inhibitor) | 2.5 µl | 2.5 µl | 2.5 µl | |
| 1x PARP buffer | 10 µl | - | - | - |
| TNKS1 (~ 2 ng/µl) | – | 10 µl | 10 µl | 10 µl |
| Total | 25 µl | 25 µl | 25 µl | 25 µl |

- 4) Add 2.5 µl of inhibitor solution to each well designated “Test Inhibitor”. For the “Positive Control”, “Substrate Control”, and “Blank”, add 2.5 µl of the same solution without inhibitor (inhibitor buffer).
- 5) Add 10 µl of 1x PARP buffer to the well designated “Blank”.
- 6) Initiate the reactions by adding 10 µl of diluted TNKS1 prepared as described above. Incubate the reactions for 1 hour at room temperature.
- 7) Wash the plate 3 times with 100 µl PBST per well.

Step 3: Detection

- 1) Dilute **Streptavidin-HRP** 1:50 in Blocking buffer.
- 2) Add 25 µl of diluted **Streptavidin-HRP** to each well. Incubate for 30 minutes at room temperature.
- 3) Wash three times with 100 µl PBST buffer as above.
- 4) Just before use, mix on ice 25 µl **HRP chemiluminescent substrate A** and 25 µl **HRP chemiluminescent substrate B** and add 50 µl per well. Discard any unused chemiluminescent reagent after use.
- 5) Incubate for 5 minutes at room temperature then read sample in a luminometer or microtiter-plate reader capable of reading chemiluminescence.


OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com

Example of Assay Results:



Inhibition of TNKS1 enzyme (Cat. # 80504) with XAV939 (Cat. # 27100), measured using the TNKS1 Histone Ribosylation Assay Kit (Biotin-labeled NAD⁺), BPS Bioscience (Cat. # 80579). Luminescence was measured using a Bio-Tek fluorescent microplate reader. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

RELATED PRODUCTS:

| <u>Product</u> | <u>Cat. #</u> | <u>Size</u> |
|---|---------------|-------------|
| TNKS1 Histone Ribosylation Assay Kit (Antibody Detection) | 80574 | 96 rxns. |
| TNKS2 Histone Ribosylation Assay Kit (Antibody Detection) | 80576 | 96 rxns. |
| TNKS2 Histone Ribosylation Assay Kit (Biotin-labeled NAD ⁺) | 80578 | 96 rxns. |
| TNKS1 (PARP5A) Enzyme | 80504 | 10 µg |
| TNKS2 (PARP5B), (667-end) Enzyme | 80505 | 10 µg |
| TNKS2 (PARP5B), (849-end) Enzyme | 80515 | 10 µg |
| PARP5b (TNKS2) Assay Kit | 80576 | 96 rxns. |
| PARP6 Assay Kit | 80556 | 32 rxns. |
| PARP1 Enzyme | 80501 | 10 µg |
| PARP2 Enzyme | 80502 | 10 µg |
| PARP3 Enzyme | 80503 | 10 µg |
| PARP6 Enzyme | 80506 | 10 µg |

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

To place your order, please contact us by Phone **1.858.829.3082** Fax **1.858.481.8694**

Or you can Email us at: info@bpsbioscience.com

Please visit our website at: www.bpsbioscience.com