




Rabbit antibody to ATG4C (APG4C): whole serum

Catalogue No.:	R-156-100
Description:	FUNCTION: Cysteine protease required for autophagy, which cleaves the C-terminal part of either MAP1LC3, GABARAPL2 or GABARAP, allowing the liberation of form I. A subpopulation of form I is subsequently converted to a smaller form (form II). Form II, with a revealed C-terminal glycine, is considered to be the phosphatidylethanolamine (PE)-conjugated form, and has the capacity for the binding to autophagosomes. ENZYME REGULATION: Inhibited by N-ethylmaleimide. SUBCELLULAR LOCATION: Cytoplasm (Probable). SIMILARITY: Belongs to the peptidase C54 family.
Batch No.:	See product label
Unit size:	100 µl
Antigen:	A synthetic peptide (CTVYNSDVIDKQ) corresponding to the C-terminal of human ATG4C protein conjugated to Blue Carrier Protein has been used as the immunogen. The peptide is homologous with the corresponding sequence derived from ATG4C protein in mouse and rat.
Other Names:	Cysteine protease ATG4C; Autophagy-related protein 4 homolog C; Autophagin-3; Autophagy-related cysteine endopeptidase 3; AUT-like 3 cysteine endopeptidase; ATG4C; APG4C; AUTL3;
Accession:	ATG4 C_HUMAN ATG4 C_MOUSE ATG4 C_RAT
Produced in:	Rabbit
Purity:	Whole serum
Applications:	IHC, immunofluorescence and WB. A dilution of 1:100 to 1:2000 dilution is recommended for these applications. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	IHC, ELISA and WB confirmed the specificity for ATG4C.
Cross-reactivity:	Human, rat. Other species not yet tested.
Form:	Lyophilised
Reconstitution:	Reconstitute in 100 µl of sterile water. Centrifuge to remove any insoluble material.
Storage:	After reconstitution keep aliquots at -20°C for a higher stability, and at 4°C with an appropriate antibacterial agent. Glycerol (1:1) may be added for an additional stability. Avoid repetitive freeze/thaw cycles.
Expiry Date:	Six months after purchase
References:	1. Marino G, et al. J. Biol. Chem. 278:3671-3678(2003). 2. Ota T, et al. Nat. Genet. 36:40-45(2004).

FOR RESEARCH USE ONLY



Rabbit antibody to ATG4C (APG4C): whole serum



Confocal microscopy on immunofluorescently detected APG4C in HL60 cell line using Rabbit antibody to ATG4C (APG4C): whole serum (R-156-100) at a dilution of 1: 100, incubated for 1 h at room temperature. The cells stained for APG4C appear in red. The cells were then counter stained with Hoechst Dye (blue colour). Here, the merged picture is presented. Note: More IHC analysis is underway. Please enquire.

biosensis

FOR RESEARCH USE ONLY