

Rabbit polyclonal antibody to Nicastrin, N-terminal domain: IgG

Catalogue No.: R-1686-500

Description: Nicastrin, a type 1 membrane glycoprotein, is an essential component of the gamma secretase

complex which is critical for the cleavage of the amyloid precursor protein and other membrane proteins. Nicastrin is widely expressed in different tissue types. This antibody detects all

processed forms of Nicastrin.

Batch No.: See vial label

Unit size: 500 ug

Antigen: A synthetic peptide (RGNSVERKIYIPL-C) corresponding to human Nicastrin [32-44] in the

N-terminal region conjugated via additional C-terminal Cys to Diphtheria toxoid.

Antibody Type: Polyclonal Produced in: Rabbit

Applications: WB and IP. Suggested concentration of 3-10 ug/mL is recommended for WB. Human or mouse

brain samples commonly prepared with reducing (50mM DTT), urea (2.3 M), SDS (1.5%) in 62.5 mM Tris-HCL pH 6.8 sample buffer (without boiling) heating to 50 C for 15 min. Unprocessed full length human Nicastrin is 709 amino acids, however this protein contains an N-terminal signal peptide which is considered to undergo cleavage during processing and transit to the cell plasma membrane, in addition the protein undergoes glycosylation to produce a glycoprotein of about 145 kDa apparent MW by SDS PAGE. Biosensis recommends that the

optimal working dilution should be determined by the end user.

Specificity: Confirmed by WB using peptide absorption.

Species Against: Confirmed by WB using peptide absorption.

Form: Lyophilized from PBS, pH 7.4. Contains no preservative.

Reconstitution: Reconstitute in 500 µL of sterile water. Centrifuge to remove any insoluble material.


Storage: Short term storage at 2-8°C for one week. At -20°C as an undiluted liquid for up to 12 months.

Expiry Date: 12 months after purchase

References: 1. Ilaya, N.T., Evin, G., Masters, C.L., and Culvenor, J.G. (2004) Nicastrin expression in mouse

peripheral tissues is not co-ordinated with Presenilin and is high in muscle. J. Neurochem. 91,

230-237.



FOR RESEARCH USE ONLY