
Product Information



Catalog No. 10007208

Sold under license from Lipid Maps




Formal Name: 9α-hydroxy-11,15-dioxo-prost-5Z-en-1-oic

Synonym: 13,14-dihydro-15-keto PGD,

MF: $C_{20}H_{32}O_5$ FW: 352.5 **Purity:** ≥98%

Stability: ≥1 year at -80°C

Supplied as: A solution in methyl acetate



Laboratory Procedures

For long term storage, we suggest that 13,14-dihydro-15-keto prostaglandin D₂ (13,14-dihydro-15-keto PGD₂) be stored as supplied at -80°C. It should be stable for at least one year.

13,14-dihydro-15-keto PGD2 is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, or dimethyl formamide purged with an inert gas can be used. The solubility of 13,14-dihydro-15-keto PGD2 in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free aqueous solution of 13,14-dihydro-15-keto PGD₂ is needed, it can be prepared by evaporating the methyl acetate and directly dissolving the neat oil in aqueous buffers. The solubility of 13,14-dihydro-15-keto PGD₂ in PBS (pH 7.2) is approximately 2.5 mg/ml. Store aqueous solutions of 13,14-dihydro-15-keto PGD₂ on ice and use within 12 hours of preparation.

13,14-dihydro-15-keto PGD2 is a metabolite of PGD2 which is formed through the 15-hydroxy PGDH pathway. 13,14-dihydro-15-keto PGD₂ was recently identified as a selective agonist for the CRTH2/DP₂ receptor. It also inhibits ion flux in a canine colonic mucosa preparation. In humans, 13,14-dihydro-15-keto PGD, is further metabolized to give 11β-hydroxy compounds which have also undergone β-oxidation of one or both side chains. Virtually no 13,14-dihydro-15-keto PGD₂ survives intact in the urine.^{3,4}

References

- 1. Hirai, H., Tanaka, K., Yoshie, O., et al. Prostaglandin D₂ selectivity induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH2. J. Exp. Med. 193(2), 255-261 (2001).
- Rangachari, P.K. and Betti, P.-A. Biological activity of metabolites of PGD2 on canine proximal colon. Am. J. Physiol. 264, G886-G894 (1993).
- 3. Liston, T.E. and Roberts, L.J., II Metabolic fate of radiolabeled prostaglandin D₂ in a normal human male volunteer. J. Biol. Chem. 260, 13172-13180 (1985).
- Morrow, J.D., Prakash, C., Awad, J.A., et al. Quantification of the major urinary metabolite of prostaglandin D₂ by a stable isotope dilution mass spectrometric assay. Anal. Biochem. 193, 142-148 (1991).

Related Products

Prostaglandin Metabolite HPLC Mixture - Cat. No. 10005 • Prostaglandin D2 - Cat. No. 12010 • 11-deoxy-11-methylene-15-keto Prostaglandin D2 - Cat. No. 12415 • 13,14-dihydro-15-keto Prostaglandin D2 - Cat. No. 12610 • 15-hydroxy Prostaglandin Dehydrogenase Polyclonal Antibody - Cat. No. 160615 • 13,14-dihydro-15-keto Prostaglandin D2-d4

WARNING: This product is for laboratory research only: not for administration to humans. Not for human or veterinary DIAGNOSTIC OR THERAPEUTIC USE.

MATERIAL SAFETY DATA

This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, or inhale. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Before use, the user must review the complete Material Safety Data Sheet, which has been sent under separate cover to the MSDS supervisor at your institution.

Cayman Chemical Company makes no warranty or guarantee of any kind, whether written or oral, expressed or implied, including without limitation, any warranty of fitness for a particular prose, suitability and merchantability, which extends beyond the description of the chemicals hereof. Cayman warrants only to the original customer that the material will meet our specifications

at the time of delivery.

Cayman will carry out its delivery obligations with due care and skill. Thus, in no event will Cayman have any obligation or liability, whether in tort (including negligence) or in contract, for any direct, inclidental or consequential damages, even if Cayman is informed about their possible existence.

This limitation of liability does not apply in the case of intentional acts or negligence of Cayman, is directors or its employees.

Buyer's exclusive remedy and Cayman's sole liability hereunder shall be limited to a refund of the purchase price, or at Cayman's option, the replacement, at no cost to Buyer, of all material that

tools not mere our specimentous.

Said refund or replacement is conditioned on Buyer giving written notice to Cayman within thirty (30) days after arrival of the material at its destination. Failure of Buyer to give said notice within thirty (30) days shall constitute a waiver by Buyer of all claims hereunder with respect to said material.

For further details, please refer to our Warranty and Limitation of Remedy located on our website and in our catalog.

Copyright Cayman Chemical Company, 11/22/2008

Cayman Chemical

Mailing address

1180 E. Ellsworth Road Ann Arbor, MI 48108 USA

(800) 364-9897 (734) 971-3335

(734) 971-3640

custserv@caymanchem.com

www.caymanchem.com