

Mouse Anti-Chicken Ig λ Monoclonal Antibody, FITC Conjugated

Mouse, Monoclonal (Immunoglobulin λ)

Cat. No. DMAB4659

Lot. No. (See product label)

PRODUCT INFORMATION

Product Overview: Mouse Monoclonal Antibody to Chicken Immunoglobulin λ light chains specific

Clone: M-2

Ig Isotype: Mouse IgG_{1k}

Format: Fluorescein (FITC) Conjugate

Quality: 0.5 mg

Specificity: λ light chains of chicken immunoglobulin

Applications: Immunofluorescence staining; Immunohistochemistry (acetone-fixed, frozen sections); ELISA

Characterization: To ensure lot-to-lot consistency, each batch of monoclonal antibody is tested to conform to characteristics of a standard reference reagent using SDS-PAGE analysis and ELISA.

Working Dilutions:

Flow Cytometry: $\leq 1 \mu\text{g}/10^6$ cells;

Other Applications: Since applications vary, each investigator should determine the optimum working dilutions of the product that is appropriate for their specific needs.

Handling And Storage: The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C. Protect conjugated forms from light. Each reagent is stable for the period shown on the bottle label if stored as directed.

Warning: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

BACKGROUND

Introduction: Lambda light chain is the smaller of the two types of polypeptide chains in immunoglobulins, consisting of an antigen-binding segment with a variable amino acid sequence, and a constant region with a relatively unchanging amino acid sequence. The lambda light chain gene is located on chromosome 22. The immunoglobulin heavy chains (G, A, M, D, or E) and light chains (kappa or lambda) are produced separately by the plasma cells in the bone marrow. The kappa and lambda molecules are attached to the heavy chains and whole immunoglobulins are assembled, and then conveyed to the surface of the plasma cell. Small amounts of free kappa and lambda light chains are found in the serum of healthy individuals since they are produced in excess. An accurate measurement of serum levels of free light chains in the presence of a great excess of bound light chains has been quite tricky for many years due to their identical structure. A number of diseases affect the production of free light chains by the plasma cells in the bone marrow, causing abnormal levels in the serum, urine or cerebrospinal fluid. Lambda along with kappa light chains can be used as a tumor marker in urine.

Keywords: Ig; Immunoglobulin ; Ig γ light chain; Immunoglobulin γ light chain; Ig lambda light chain; Immunoglobulin lambda light chain

REFERENCES

1. Weigert. M. & Riblet. R. Cold Spring Harb Symp. quant. Biol. 41. 837-846(1976).
2. Bernard, O., Hozumi, N. & Tonegawa, S. Cell 15, 1133-1144(1978)

Creative Diagnostics. All rights reserved.

45-16 Ramsey Road Shirley, NY 11967, USA
Tel: 631-624-4882 · Fax: 631-614-7828
E-mail: info@creative-diagnostics.com
www.creative-diagnostics.com