



EB06598 - Goat Anti-ABAD / HADH2 Antibody

Size: 100µg specific antibody in 200µl



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

Target Protein

Principal Names: HADH2 antibody, ABAD antibody, ERAB antibody, MHBD antibody, HSD17B10 antibody, 17b-HSD10 antibody, hydroxyacyl-Coenzyme A dehydrogenase, type II antibody, type 10 17b-HSD antibody, AB-binding alcohol dehydrogenase antibody, type 10 17beta-hydroxysteroid dehydrogenase antibody

Official Symbol: HSD17B10

Accession Number(s): NP_004484.1; NP_001032900.1

Human GeneID(s): [3028](#)

Important Comments: This antibody is expected to recognise both reported isoforms.

Immunogen

Peptide with sequence CIRLDGAIRMQP, from the C Terminus of the protein sequence according to NP_004484.1; NP_001032900.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 25kDa band observed in Human Brain lysates (calculated MW of 26.9kDa according to NP_004484). Recommended concentration: 0.3-2µg/ml. A minor band of unknown identity was also consistently observed at appr 38kDa. This band was successfully blocked by incubation with the immunising peptide.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference

The goat polyclonal antibody used (in WB on Mouse tissue) as described in this paper was manufactured by us:

Tilton RG, Haidacher SJ, Lejeune WS, Zhang X, Zhao Y, Kurosky A, Brasier AR, Denner L.

Diabetes-induced changes in the renal cortical proteome assessed with two-dimensional gel electrophoresis and mass spectrometry.

Proteomics. 2007 May;7(10):1729-42.

PMID: 17436268



EB06598 staining (0.3 μ g/ml) of Human Brain lysate (RIPA buffer, 30 μ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.