

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire

UK

Enquiries:

OX25 5HD

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06659 - Goat Anti-ACHE Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ACHE antibody, YT antibody, acetylcholinesterase (YT blood group) antibody, ARACHE antibody, N-ACHE antibody, OTTHUMP00000211347 antibody,

OTTHUMP00000211349 antibody, OTTHUMP00000211356 antibody, acetylcholinesterase; apoptosis-related acetylcholinesterase antibody

Official Symbol: ACHE

Accession Number(s): NP_000656.1

Human GeneID(s): 43

Important Comments: This antibody is expected to recognise isoform NP_000656 only

(the ubiquitously expressed, hydrophillic form).

Immunogen

Peptide with sequence QFDHYSKQDRCSDL, from the C Terminus of the protein sequence according to NP_000656.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 70kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 67.8kDa according to NP_000656.1). Recommended concentration: 1-3μg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow



EB06659 (1 μ g/ml) staining of Human Cerebellum lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.