



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07069 - Goat Anti-GAPDH (Internal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GAPDH antibody, glyceraldehyde-3-phosphate dehydrogenase antibody, HGNC:4141 antibody, G3PD antibody, GAPD antibody, MGC88685 antibody, OTTHUMP00000174431 antibody, OTTHUMP00000174432 antibody, aging-associated gene 9 protein antibody, glyceraldehyde 3-phosphate dehydrogenase antibody

Official Symbol: GAPDH

Accession Number(s): NP_002037.2

Human GeneID(s): [2597](#)

Important Comments: GAPDH is constitutively expressed in almost all tissues at high levels. It is therefore a useful marker when a loading/positive control is required in western blotting.

Immunogen

Peptide with sequence C-GVNHEKYDNSLK, from the internal region of the protein sequence according to NP_002037.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 36kDa band observed in Human Liver, Testes and Tonsil lysates and in Mouse Spleen lysates (calculated MW of 36.1kDa according to NP_002037.2).

Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

Specific References

This antibody has been successfully used in the following paper:

Carr AJ, Vugler AA, Yu L, Semo M, Coffey P, Moss SE, Greenwood J.

The expression of retinal cell markers in human retinal pigment epithelial cells and their augmentation by the synthetic retinoid fenretinide.

Mol Vis. 2011;17:1701-15.

PMID: 21738400

This antibody has been successfully used in the following paper:

Carr AJ, Vugler A, Lawrence J, Chen LL, Ahmado A, Chen FK, Semo M, Gias C, da Cruz L, Moore HD, Walsh J, Coffey PJ.

Molecular characterization and functional analysis of phagocytosis by human embryonic stem cell-derived RPE cells using a novel human retinal assay.

Mol Vis. 2009;15:283-95.

PMID: 19204785

This antibody has been successfully used in the following paper:

Vugler A, Carr AJ, Lawrence J, Chen LL, Burrell K, Wright A, Lundh P, Semo M, Ahmado A, Gias C, da Cruz L, Moore H, Andrews P, Walsh J, Coffey P.

Elucidating the phenomenon of HESC-derived RPE: anatomy of cell genesis, expansion and retinal transplantation.

Exp Neurol. 2008 Dec;214(2):347-61.

PMID: 18926821

The goat polyclonal antibody used (in WB on Dog heart) as described in this paper was manufactured by us:

Chakir K, Daya SK, Aiba T, Tunin RS, Dimaano VL, Abraham TP, Jaques-Robinson KM, Lai EW, Pacak K, Zhu WZ, Xiao RP, Tomaselli GF, Kass DA.

Mechanisms of enhanced beta-adrenergic reserve from cardiac resynchronization therapy.

Circulation. 2009 Mar 10;119(9):1231-40.

PMID: 19237665


The goat polyclonal antibody used (in WB on Human lymphocytes (PBMC)) as described in this paper was manufactured by us:

Schillace RV, Miller CL, Pisenti N, Grotzke JE, Swarbrick GM, Lewinsohn DM, Carr DW.

A-kinase anchoring in dendritic cells is required for antigen presentation.

PLoS One. 2009;4(3):e4807.

PMID: 19277197



EB07069 (0.1 μ g/ml) staining of Human Tonsil Lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.