



EB07373 - Goat Anti-APBB1 / FE65 Antibody

Size: 100µg specific antibody in 200µl



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

Target Protein

Principal Names: APBB1 antibody, FE65 antibody, amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65) antibody, MGC:9072 antibody, RIR antibody, adaptor protein FE65a2 antibody, amyloid beta A4 precursor protein-binding, family B, member 1 antibody, stat-like protein antibody

Official Symbol: APBB1

Accession Number(s): NP_001155.1; NP_663722.1

Human GeneID(s): [322](#)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_001155.1 and NP_663722.1).

Immunogen

Peptide with sequence C-GSLKPKRLGAHTP, from the C Terminus of the protein sequence according to NP_001155.1; NP_663722.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 100kDa band observed in lysates of NIH/3T3 and HeLa (calculated MW of 77.2kDa according to NP_001155.1 and 77.0kDa according to NP_663722.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Zambrano et al, J Biol Chem. 1998 Aug 7;273(32):20128-33.; PMID: 9685356). Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human



EB07373 (0.1 μ g/ml) staining of NIH/3T3 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.