

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09039 - Goat Anti-MEIS1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MEIS1 antibody, Meis homeobox 1 antibody, MGC43380 antibody,

Meis1, myeloid ecotropic viral integration site 1 homolog antibody,

WUGSC:H_NH0444B04.1 antibody, leukemogenic homolog protein antibody

Official Symbol: MEIS1

Accession Number(s): NP_002389.1

Human GeneID(s): 4211

Non-Human GenelD(s): 17268 (mouse), 686117 (rat)

Immunogen

Peptide with sequence C-SEDITRSANLTDQ, from the internal region of the protein sequence according to NP_002389.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx. 40kDa band observed in Human Brain (Cerebral Cortex) lysates (calculated MW of 43.0kDa according to NP_002389.1). Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB09039 (0.1 μ g/ml) staining of Human Cerebral Cortex lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.