



UK Office

Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore
405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09983 - Goat Anti-CLCN2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: chloride channel 2 antibody, CIC-2 antibody, CLC2 antibody, ECA2 antibody, ECA3 antibody, EGI11 antibody, EGI3 antibody, EGMA antibody, EJM6 antibody, EJM8 antibody, Epilepsy, idiopathic generalized, susceptibility to antibody, CLCN2 antibody

Official Symbol: CLCN2

Accession Number(s): NP_004357.3

Human GeneID(s): [1181](#)

Non-Human GeneID(s): 12724 (mouse), 29232 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_004357.3; NP_001164558.1; NP_001164559.1; NP_001164560.1).

Immunogen

Peptide with sequence QQLDEPVNFSDCK, from the internal region of the protein sequence according to NP_004357.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 100kDa band observed in lysates of cell line HeLa (calculated MW of 98.5kDa according to NP_004357.3). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow, Rabbit



EB09983 (0.3 μ g/ml) staining of HeLa lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.