

CREATING NOVEL ANTIBODIES

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire **OX25 5HD** 

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona. CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10037 - Goat Anti-OCT4 / POU5F1 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: Oct3 antibody, Oct-3 antibody, Oct3/4 antibody, Oct-3/4 antibody, Oct-4 antibody, Oct-4 antibody, Otf3 antibody, Otf-3 antibody, Otf3g antibody, Otf3-rs7 antibody, Otf4 antibody, Otf-4 antibody, POU domain, class 5, transcription factor 1 antibody,

Pou5f1 antibody

Official Symbol: POU5F1

Accession Number(s): NP\_038661.2 Non-Human GeneID(s): 18999 (mouse)

## **Immunogen**

Peptide with sequence C-DRPNAVKLEKVEP, from the internal region of the protein sequence according to NP\_038661.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 40kDa band observed in lysates of cell line NIH3T3 (calculated MW of 38.2kDa according to NP\_038661.2). Recommended concentration: 1-3µg/ml.

# **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Mouse, Rat



EB10037 (1µg/ml) staining of NIH3T3 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.