Shikhar Biotech Pvt Ltd Khumaltar, Lalitpur P.O.Box 21608 Kathmandu, Nepal

Phone: +977-1-5545788 Email: Info@shikharbiotech.com



Product Datasheet

Shi-fix[™] coverslips

#SB-Shifix25 / SB-Shifix25 PLUS /SB-Shifix50 / SB-Shifix50PLUS

Description Glass coverslips coated on both sides with Shi-fix[™] biomaterial that

enhances binding of non adherent animal cells for imaging studies. Shi-fix $^{\text{\tiny{TM}}}$ cover-slips allow suspension cells to be layered on them for imaging

or cultured as a monolayer. These cover-slips are suitable for immunofluorescence and confocal microscopy. Cells can also be

stimulated while they are attached on the cover-slips.

Dimensions Shi-fix[™] : #1.5, ø 12mm

Shi-fix™ PLUS: #1.5, ø 15mm

Packing size SB-Shifix25 (pack of 25) SB-Shifix50 (pack of 50) SB-Shifix50PLUS (pack of

50).

Intended use This product is intended for research use only.

Storage and handling The product can be shipped at ambient temperature. Unopened product

can be stored at room temperature but needs to be kept at 4C after

opening the sealed packaging.

The shelf life is 6 months at room temperature and 1 year at 4C.

Procedure for Use 1. Wash cells with PBS and resuspend 2-5 million cells per ml in PBS

2. Place the Shi-fix[™] cover-slips on a 12-well ELISA plate for ease of handling. Add the cells directly to Shi-fix[™] cover-slips.(0.2-0.3 million cells

nandling. Add the cells directly to Shi-fix'" cover-slips.(0.2-0.3 million cells

per coverslip).

3. Let the cells settle and attach for 30 mins. Attachment for some cell types may improve by incubating longer but do not exceed 1 hour. Wash gently with 2ml PBS to remove unbound cells (*Do not pipet PBS straight*

onto the coverslips. Add PBS to the sides of the plate well followed by

gentle rocking for 3-5 mins).

4. Check for cell attachment on a microscope. If further culturing is needed, add 1ml medium otherwise proceed to fixation, permeabilization

and immunostaining.

Safety This product is for use by personnel trained in mammalian cell culture

techniques. Lab coat and gloves must be worn. Coverslips must be discarded after use according to the applicable biological waste

regulations of the institute or department.