

Myc Tag Antibody

Catalog Number: E1T506-1, E1T506-2

Amount: 50μg/50μl, 100μg/100μl Form of Antibody: Mouse Monoclonal (IgG1)

1mg/ml in PBS, pH 7.4 with 0.05% sodium azide. Frozen liquid.

Support: service@enogene.com

Clone Myc.A7

Storage/Stability: Stable for one year at -20°C from date of shipment. For maximum recovery of product,

centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid

repeated freezing and thawing. Aliquot will be stable at 4°C for 3 months.

A synthetic peptide corresponding to residues 410-419 (EQKLISEEDL) coupled to KLH. Immunogen:

Specificity/Sensitivity: Myc-tag Monoclonal Antibody. Recognizes over-expressed proteins containing Myc epitope

tag fused to either amino- or carboxy-termini of targeted proteins in transfected mammalian


Reactivity: Human, Mouse, Rat, others

Applications: Optimal working dilutions should be determined experimentally by the investigator. Prepare

> working dilution immediately before use. Suggested starting dilutions are as follows: Western Blot at dilutions of 1:1000-4000. Immunoprecipitation and immunostaining at

dilutions of 1:200-400.

Watanabe S, et al. (2006) Mol Reprod Dev; 73(11):1345-1352. References:



Western Blot analysis of Myc-tagged fusion construct under different transfection conditions of transfection reagents and amount of fusion DNA construct.

For Research Use Only