

CD4 FITC - CD28 PE. Mouse Bi-Test™ Reagents (FITC/RPE)

COMMENTS

Identification of CD4 on human helper/inducer T cells expressing the 60,000 M.W. surface antigen (HLA class II reactive). CD4 is present in low density on monocytes. Anti-human CD28 binds the 44kDa MW cell surface protein on the surface of most T cells. CD28 acts as the ligand for the B7/BB-1 molecule on the surface of activated B cells. B7/BB-1 costimulates T cells through CD28, along with CD2 and CD3. CD28 antigen is a disulfide-linked homodimeric glycoprotein. The CD28 antigen is present on approximately 60%-80% of lymphocytes (95% of CD4 and 50% of CD8 lymphocytes). CD28 regulates the expression of cytokines by T cells, not only IL-2, but also IL-1 alpha and CSF-1, usually synthesized by accessory cells. CD28 functions as a cell adhesion molecule (CAM) for certain T cell subsets.

CONCENTRATION

See vial for concentration

SHIP CONDITIONS

Room Temperature

STORAGE CUSTOMER

Product should be stored at 4-8°C. DO NOT FREEZE

STABILITY

Reagents are stable for the period shown on the vial label when stored properly

Use

PBMC: Add10 μ l of MAB/10^6 PBMC in 100 μ l PBS. Mix gently and incubate for 15 minutes at 2° to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add10 μ l of MAB/100 μ l of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

ORDERING INFORMATION

CATALOG NUMBER

0428S

SIZE

50 Tests

FORM

Bi-Test (FITC/RPE) Reagent

HOST/CLONE

Mouse

FORMULATION

Provided as sterile filtered solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein

ISOTYPE

IgG1 (F)/IgG1 (PE)

APPLICATIONS

Flow Cytometry