

## CD45RA FITC. Mouse Single-Color Reagent BHKG-2, Human

## COMMENTS

Anti Human Leukocyte Common Antigen FITC

**CONCENTRATION** See vial for concentration

SHIP CONDITIONS Room Temperature STORAGE CUSTOMER

Product should be stored at 4-8°C. DO NOT FREEZE

STABILITY

Reagents are stable for the period shown on the vial label when stored properly

Use

Add 10 µl of MAB / 100 µl of 10^6 PBMC in PBS. Mix Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. Add 10 µl of MAB / 100 µl of Whole Blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse Whole Blood and analyze or fix with 0.5% v/v paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

Last Modified 3/22/2012

For research use only. Not for use in human diagnostics or therapeutics.

Page 1 of 2 Cat. No. 045R2

Exalpha Biologicals, Inc. 2 Shaker Road, Bldg. B101 Shirley, MA 01464 Tel: 800.395.1137 Fax: 866.924.5100 www.exalpha.com info@exalpha.com

045R2 Size 100 Tests FORM FITC HOST/CLONE Mouse Clone BHKG-2 FORMULATION Provided as sterile filtered solution in phosphate buffered saline with 0.08%

**ORDERING INFORMATION** 

CATALOG NUMBER

phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein **ISOTYPE** IgG1

APPLICATIONS Flow Cytometry