



11-137-C025

Monoclonal Antibody to CD49c / Integrin alpha 3 Purified Antibody (0.025 mg)

Clone:	ASC-1
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody ASC-1 recognizes CD49c (integrin alpha 3), a transmembrane glycoprotein composed of disulfide linked 125 kDa and 30 kDa chains, and expressed on adherent cell lines and to a lesser extent on T and B cells and monocytes. HLDA VI; WS Code A002
Regulatory Status:	RUO
Immunogen:	Human SSC-9 cell line (squamous cell carcinoma)
Species Reactivity:	Human
Application:	Flow Cytometry Immunoprecipitation Western Blotting Immunohistochemistry (frozen sections)
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix components (fibronectin, laminin 1, laminin 5, entactin, and collagen). It is expressed on adherent cells, mainly on fibroblasts, epithelial cells and endothelial cells.
References:	*Pattaramalai S, Skubitz KM, Skubitz AP: A novel recognition site on laminin for the alpha 3 beta 1 integrin. <i>Exp Cell Res.</i> 1996 Feb 1;222(2):281-90. *Skubitz AP, Bast RC, Wayner EA, Letourneau PC, Wilke MS: Expression of alpha 6 and beta 4 integrins in serous ovarian carcinoma correlates with expression of the basement membrane protein laminin. <i>Am J Pathol.</i> 1996 May;148(5):1445-61.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz