



11-139-C025

Monoclonal Antibody to CD89 Purified Antibody (0.025 mg)

Clone:	A59
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody A59 recognizes CD89, a 55-100 kDa glycoprotein serving as a receptor for IgA and expressed mainly on granulocytes, monocytes and macrophages. HLDA WS Code V MR30
Regulatory Status:	RUO
Immunogen:	Ag8.653 myeloma cells
Species Reactivity:	Human, Non-Human Primates
Application:	Flow Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89 is expressed on granulocytes, monocytes, macrophages, dendritic cells and myeloid cell lines. Its expression is upregulated in presence of IgA immune complexes, stimulators (such as LPS, PMA), TNF alpha, IL1 beta or GM-CSF, and it is downregulated in presence of TGF beta and suramin. Binding of IgA-opsonized targets to CD89 leads to phagocytic and cytotoxic processes of the immunologic defense.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Schenk M, Bouchon A, Seibold F, Mueller C: TREM-1--expressing intestinal macrophages crucially amplify chronic inflammation in experimental colitis and inflammatory bowel diseases. *J Clin Invest.* 2007 Oct;117(10):3097-106.
- *Barratt J, Greer MR, Pawluczyk IZ, Allen AC, Bailey EM, Buck KS, Feehally J: Identification of a novel Fc α receptor expressed by human mesangial cells. *Kidney Int.* 2000 May;57(5):1936-48.
- *van Egmond M, van Spriël AB, Vermeulen H, Huls G, van Garderen E, van de Winkel JG: Enhancement of polymorphonuclear cell-mediated tumor cell killing on simultaneous engagement of fc γ RI (CD64) and fc α RI (CD89). *Cancer Res.* 2001 May 15;61(10):4055-60.
- *Honorio-França AC, Launay P, Carneiro-Sampaio MM, Monteiro RC: Colostral neutrophils express Fc alpha receptors (CD89) lacking gamma chain association and mediate noninflammatory properties of secretory IgA. *J Leukoc Biol.* 2001 Feb;69(2):289-96.
- *Wu J, Ji C, Xie F, Langefeld CD, Qian K, Gibson AW, Edberg JC, Kimberly RP: Fc α RI (CD89) alleles determine the proinflammatory potential of serum IgA. *J Immunol.* 2007 Mar 15;178(6):3973-82.
- *Rogers KA, Scinicariello F, Attanasio R: Identification and characterization of macaque CD89 (immunoglobulin A Fc receptor). *Immunology.* 2004 Oct;113(2):178-86.
- *van Dijk TB, Bracke M, Caldenhoven E, Raaijmakers JA, Lammers JW, Koenderman L, de Groot RP: Cloning and characterization of Fc alpha Rb, a novel Fc alpha receptor (CD89) isoform expressed in eosinophils and neutrophils. *Blood.* 1996 Dec 1;88(11):4229-38.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz