



Antibodies

11-155-C100

## Monoclonal Antibody to Ki-67 Purified Antibody (0.1 mg)

<b>Clone:</b>	Ki-67
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody Ki-67 recognizes Ki-67 antigen, a non-histone nuclear protein expressed exclusively in proliferating cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Nuclei of the Hodgkin lymphoma cell line L428
<b>Species Reactivity:</b>	Human, Bovine
<b>Application:</b>	Flow Cytometry Intracellular staining Western Blotting Immunohistochemistry Immunocytochemistry Application note: paraformaldehyde fixation; recommended antibody concentration 1 µg/ml
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	Ki-67 is a highly protease-sensitive nuclear protein expressed in two isoforms (345 kDa and 395 kDa), both of which are identified by the antibody clone Ki-67. The Ki-67 antigen is essential for cell proliferation and its expression is restricted to the cycling cells. It is detected in G1, S, G2 and M phase, whereas it is absent in cells which are in G0 phase and it is not associated with DNA repair processes. Ki-67 thus represents an important tool for detection of proliferating cells, which is of great importance in tumor diagnostics and is commonly used as a prognostic factor in cancer studies.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*Gerdes J, Schwab U, Lemke H, Stein H: Production of a mouse monoclonal antibody reactive with a human nuclear antigen associated with cell proliferation. *Int J Cancer*. 1983 Jan 15;31(1):13-20.
  - \*Gerdes J: Ki-67 and other proliferation markers useful for immunohistological diagnostic and prognostic evaluations in human malignancies. *Semin Cancer Biol*. 1990 Jun;1(3):199-206.
  - \*Gerdes J, Lemke H, Baisch H, Wacker HH, Schwab U, Stein H: Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. *J Immunol*. 1984 Oct;133(4):1710-5.
  - \*Duchrow M, Schlüter C, Key G, Kubbutat MH, Wohlenberg C, Flad HD, Gerdes J: Cell proliferation-associated nuclear antigen defined by antibody Ki-67: a new kind of cell cycle-maintaining proteins. *Arch Immunol Ther Exp (Warsz)*. 1995;43(2):117-21.
  - \*Schlüter C, Duchrow M, Wohlenberg C, Becker MH, Key G, Flad HD, Gerdes J: The cell proliferation-associated antigen of antibody Ki-67: a very large, ubiquitous nuclear protein with numerous repeated elements, representing a new kind of cell cycle-maintaining proteins. *J Cell Biol*. 1993 Nov;123(3):513-22.
  - \*And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)