

11-165-C100

Monoclonal Antibody to CD165 Purified Antibody (0.1 mg)

Clone:	SN2
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody SN2, also known as SN2 N6-D11, recognizes CD165, an approximately 37-42 kDa transmembrane glycoprotein expressed mainly on leukemic T cells, double positive and double negative thymocytes (CD4-CD8-, CD4+CD8+), and platelets. HLDA VI: WS Code N-L099
Regulatory Status:	RUO
Immunogen:	CD165 purified from human Molt-4 cell line
Species Reactivity:	Human
Application:	Flow Cytometry Immunoprecipitation Western Blotting Immunohistochemistry (frozen sections) Application note: acetone fixation Immunocytochemistry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified from cell culture supernatant by protein-A affinity chromatography.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD165 is a poorly characterized transmembrane protein highly expressed on platelets and many leukemic T cell lines. At lower level it is expressed on a proportion of circulating T cells and monocytes, on thymic epithelium, fibroblasts, epidermal keratinocytes, pancreatic islet cells, and some neurons. It might have a role in adhesion between thymocytes and thymic epithelial cells and it can be used as a marker for tumor progression.
References:	*Seon BK, Negoro S, Barcos MP, Tebbi CK, Chervinsky D, Fukukawa T: Monoclonal antibody SN2 defining a human T cell leukemia-associated cell surface glycoprotein. J Immunol. 1984 Apr;132(4):2089-95. *Seon BK: Specific killing of human T-leukemia cells by immunotoxins prepared with ricin A chain and monoclonal anti-human T-cell leukemia antibodies. Cancer Res. 1984 Jan;44(1):259-64.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz