



11-178-C100

Monoclonal Antibody to CD273 / PD-L2 Purified Antibody (0.1 mg)

Clone:	24F.10C12
Isotype:	Mouse IgG2a
Specificity:	The mouse monoclonal antibody 24F.10C12 recognizes CD273 / PD-L2 (also known as B7-DC), a 25 kDa type I transmembrane protein expressed by dendritic cells, activated monocytes and T cells, heart, first trimester placenta, lung and liver, as well as in Hodgkin's lymphoma cells and primary mediastinal B cell lymphoma (PMBL).
Regulatory Status:	RUO
Immunogen:	human CD273
Species Reactivity:	Human
Application:	Flow Cytometry Immunohistochemistry Functional Application blocking
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD273 / PD-L2 (programmed death ligand-1), also known as B7-DC, is a member of the B7 family of regulatory proteins. It costimulates the proliferation of T cells, and mediates IFN gamma production. Ligation of CD273 on dendritic cells enhances dendritic cell activation and T cell responses. When interacting with CD279, it can act as a coinhibitor of the T cell function. CD273 expression is a useful marker to distinguish primary mediastinal B cell lymphoma from other diffuse large B cell lymphomas.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Green MR, Monti S, Rodig SJ, Juszczynski P, Currie T, O'Donnell E, Chapuy B, Takeyama K, Neuberg D, Golub TR, Kutok JL, Shipp MA: Integrative analysis reveals selective 9p24.1 amplification, increased PD-1 ligand expression, and further induction via JAK2 in nodular sclerosing Hodgkin lymphoma and primary mediastinal large B-cell lymphoma. *Blood*. 2010 Oct 28;116(17):3268-77.
 - *Rodríguez-García M, Porichis F, de Jong OG, Levi K, Diefenbach TJ, Lifson JD, Freeman GJ, Walker BD, Kaufmann DE, Kavanagh DG: Expression of PD-L1 and PD-L2 on human macrophages is up-regulated by HIV-1 and differentially modulated by IL-10. *J Leukoc Biol*. 2011 Apr;89(4):507-15.
 - *Semnani RT, Mahapatra L, Moore V, Sanprasert V, Nutman TB: Functional and phenotypic characteristics of alternative activation induced in human monocytes by interleukin-4 or the parasitic nematode *Brugia malayi*. *Infect Immun*. 2011 Oct;79(10):3957-65.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz