



11-297-C025

Monoclonal Antibody to CD34 Purified Antibody (0.025 mg)

Clone:	4H11[APG]
Isotype:	Mouse IgG1
Specificity:	The antibody 4H11[APG] reacts with Class III epitope on CD34 (Mucosialin), a 110-115 kDa monomeric transmembrane phosphoglycoprotein expressed on hematopoietic progenitors cells and on the most pluripotential stem cells; it is gradually lost on progenitor cells. The antibody 4H11[APG] completely blocks binding of Class II antibody QBEnd10 and Class III antibodies BIRMA K3 and 8G12 on KG1a cell line. HLDA VI; WS Code M MA58
Regulatory Status:	RUO
Immunogen:	Permanent human cell line derived from peripheral leucocytes of a patient suffering from chronic myeloid leukaemia.
Species Reactivity:	Human
Application:	Flow Cytometry Recommended dilution: 2 µg/ml Western Blotting Recommended dilution: 2 µg/ml Positive control: Kg-1a human leukemia cell lysate Negative control: JURKAT human leukemia T-cell line Sample preparation: Resuspend approx. 50 mil. cells in 1 ml cold Lysis buffer (1% laurylmaltoside in 20 mM Tris/Cl, 100 mM NaCl pH 8.2, 50 mM NaF including Protease inhibitor Cocktail). Incubate 60 min on ice. Centrifuge to remove cell debris. Mix lysate with non-reducing SDS-PAGE sample buffer. Application note: Non-reducing conditions. Immunohistochemistry (paraffin sections) Recommended dilution: 10 µg/ml Positive tissue: placenta endothelium Immunocytochemistry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by precipitation and chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****Background:**

CD34 is a highly glycosylated monomeric 111-115 kDa surface protein, which is present on many stem cell populations. It is a well established stem cell marker, though its expression on human hematopoietic stem cells is reversible. CD34 probably serves as a surface receptor that undergoes receptor-mediated endocytosis and regulates adhesion, differentiation and proliferation of hematopoietic stem cells and other progenitors. CD34 expression is likely to represent a specific state of hematopoietic development that may have altered adhering properties with expanding and differentiating capabilities in both in vitro and in vivo conditions.

References:

- *Krauter J, Hartl M, Hambach L, Kohlenberg A, Gunsilius E, Ganser A, Heil G: Receptor-mediated endocytosis of CD34 on hematopoietic cells after stimulation with the monoclonal antibody anti-HPCA-1. *J Hematother Stem Cell Res.* 2001 Dec;10(6):863-71.
- *Dao MA, Arevalo J, Nolte JA: Reversibility of CD34 expression on human hematopoietic stem cells that retain the capacity for secondary reconstitution. *Blood.* 2003 Jan 1;101(1):112-8.
- *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Gulati S, Luthra PM: Three-dimensional structure prediction of the interaction of CD34 with the SH3 domain of Crk-L. *Stem Cells Dev.* 2005 Oct;14(5):470-7.
- *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Luthra PM: Hematopoietic stem cell antigen CD34: role in adhesion or homing. *Stem Cells Dev.* 2006 Jun;15(3):305-13.
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- *Elknerová K, Lacinová Z, Soucek J, Marinov I, Stöckbauer P: Growth inhibitory effect of the antibody to hematopoietic stem cell antigen CD34 in leukemic cell lines. *Neoplasma.* 2007;54(4):311-20.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.