



Antibodies

11-299-C025

## Monoclonal Antibody to CD25 Purified Antibody (0.025 mg)

<b>Clone:</b>	MEM-140
<b>Isotype:</b>	Mouse IgM
<b>Specificity:</b>	The antibody MEM-140 reacts with CD25 (Interleukin-2 receptor alpha chain), a 55 kDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4 <sup>+</sup> T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes. HLDA VI; WS Code C C-54
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	PHA-activated peripheral blood leucocytes
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Immunoprecipitation excellent for immunoisolation of CD25 <sup>+</sup> cells Flow Cytometry Recommended dilution: 1 µg/ml
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by precipitation and chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Tris buffered saline (TBS) with 15 mM sodium azide, approx. pH 8.0
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD25 (IL2Ralpha, Tac) is a ligand-binding alpha subunit of interleukin 2 receptor (IL2R). Together with beta and gamma subunit CD25 constitutes the high affinity IL2R, whereas CD25 alone serves as the low affinity IL2R. CD25 expression rapidly increases upon T cell activation. The 55 kDa CD25 molecule is enzymatically cleaved and shed from the cell surface as a soluble 45 kDa s-Tac, whose concentration in serum can be used as a marker of T cell activation. Expression of CD25 indicates the neoplastic phenotype of mast cells. Humanized anti CD25 antibodies represent a useful tool to reduce the incidence of allograft rejection as well as the severity of graft versus host reaction, and radioimmunoconjugates of anti-CD25 antibodies can be used against CD25 expressing lymphomas.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

\*Lai KN, Leung JC, Lai FM: Soluble interleukin 2 receptor release, interleukin 2 production, and interleukin 2 receptor expression in activated T-lymphocytes in vitro. *Pathology*. 1991 Jul;23(3):224-8.

\*Scheibenbogen C, Keilholz U, Richter M, Andreesen R, Hunstein W: The interleukin-2 receptor in human monocytes and macrophages: regulation of expression and release of the alpha and beta chains (p55 and p75). *Res Immunol*. 1992 Jan;143(1):33-7.

\*Morris JC, Waldmann TA: Advances in interleukin 2 receptor targeted treatment. *Ann Rheum Dis*. 2000 Nov;59 Suppl 1:i109-14.

\*Sotlar K, Horny HP, Simonitsch I, Krokowski M, Aichberger KJ, Mayerhofer M, Printz D, Fritsch G, Valent P: CD25 indicates the neoplastic phenotype of mast cells: a novel immunohistochemical marker for the diagnosis of systemic mastocytosis (SM) in routinely processed bone marrow biopsy specimens. *Am J Surg Pathol*. 2004 Oct;28(10):1319-25.

\*Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)